



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [info.bmjopen@bmj.com](mailto:info.bmjopen@bmj.com)

# BMJ Open

## Sports Injuries Aligned to Predicted Mature Height in Highly Trained Youth Athletes: Cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023284
Article Type:	Research
Date Submitted by the Author:	08-Apr-2018
Complete List of Authors:	<p>Rejeb, Abdallah; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie</p> <p>Johnson, Amanda; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center</p> <p>Farooq, Abdulaziz; Aspetar, Orthopaedic and Sports Medicine Department, Athlete Health and Performance Research</p> <p>Verrelest, Ruth ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie</p> <p>Pullinger, Samuel; ASPIRE Academy for Sports Excellence, Sports Sciences</p> <p>Vaeyens, Roel ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Department of movement and Sports Sciences</p> <p>Witvrouw, Erik; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapy</p>
Keywords:	Youth, Biological maturation, Skeletal age, Anthropometrics, Sports injury, Mature height

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Sports Injuries Aligned to Predicted Mature Height in Highly Trained Youth Athletes:**

**Cohort study**

**Authors:**

Abdallah Rejeb<sup>1 2</sup>, Amanda Johnson<sup>1</sup>, Abdulaziz Farooq<sup>1</sup>, Ruth Verrelest<sup>2</sup>, Samuel Pullinger<sup>3</sup>,  
Roel Vaeyens<sup>4</sup>, Erik Witvrouw<sup>2</sup>

**Affiliations:**

<sup>1</sup> Aspetar Orthopedic Sports Medicine and Hospital, Doha, Qatar

<sup>2</sup> Ghent University, Department of Rehabilitation Sciences and Physiotherapy, Belgium

<sup>3</sup> Aspire Academy for Sports Excellence, Doha, Qatar

<sup>4</sup> Ghent University, Department of Movement and Sports Sciences, Belgium

**Correspondence:**

Abdallah Rejeb

Address: Asian Games 2006 Road, P.O. Box 22287, Aspire Academy for Sports, Doha, Qatar

Email: [Abdallah.rajeb@aspetar.com](mailto:Abdallah.rajeb@aspetar.com)

Word count = 3445

## ABSTRACT

**Objectives:** To investigate the association of maturity status with injury incidence in highly trained Middle-Eastern youth athletes.

**Design:** Prospective cohort study

**Setting:** Four consecutive seasons ranging from 2010 to 2014 at the Aspire Academy for Sports Excellence, Qatar.

**Participants:** Male athletes (age range: 11–18 y) representing four sporting disciplines were enrolled and grouped into two categories: individual sports (athletics and fencing) and racquet sports (squash and table tennis).

**Outcome measures:** Athletes' anthropometric characteristics were assessed to calculate age at peak height velocity (APHV) and total years from PHV. Participants predicted mature heights (PMHs) were collected and categorized into four PMH quartiles. Consenting athletes had wrist and hand radiographs taken for assessment of skeletal age (SA), using Fels method. Early and late maturers were those with an SA of >1 y either older or younger than their chronological age (CA), respectively.

**Results:** For the sample (n = 67) across all sport groups, 43 (64%) athletes had one or more injuries: a total of 212 injuries, or 4.9 injuries per registered athlete. Survival analysis using Cox regression of maturity status found that early maturing athletes had a two-fold greater risk of injury over time compared to late maturers (hazard ratio [HR]; 2.04, 95% CI 1.15–3.61, P = 0.015). PMH was associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006).

Compared to the PMH for participants in the 1st quartile (<176 cm), athletes in the 4th quartile ( $\geq 184$  cm) had a higher (up to 2-fold) injury risk (HR 2.41, 95% CI 1.42–4.08, P = 0.001).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Racquet and individual sports involved a similar injury risk (HR 1.14, 95% CI 0.86–1.52, P = 0.37).

**Conclusions:** Early maturation and PMH gradient were significant predictors of injury in youths.

**Key words:** youth, biological maturation, skeletal age, anthropometrics, sports injury, mature height

**Strengths and limitations of this study**

- The first longitudinal study to assess the anthropometric characteristics and biological maturity status as injury risk factors in Middle-Eastern youth athletes.
- The participants were highly trained male adolescent athletes.
- Measurement of maturity and growth were found to be at moderate-to-high risk of bias.
- SA as an indicator of maturation, has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment protocols and the interpretation of results.

## 53 INTRODUCTION

54 The range of somatic and biological maturity in individuals of the same chronological age (CA)  
55 is large [1]. Such observations are derived from correlational and multivariate studies that  
56 compare young individuals of the same age who are at both extremes of the maturity range [2].  
57 Therefore, the assessment of maturity is an important consideration when dealing with  
58 adolescent athletes on a longitudinal basis. Further, understanding the cause of disease and  
59 injury is vital in predicting and preventing injury [3].

60 In young athletes, the demands of their chosen sport are superimposed on normal growth and  
61 maturation. A literature review revealed that there is a greater susceptibility to injury during  
62 certain periods of growth [4–6]. Indeed, the association between an increased prevalence of  
63 injuries and the adolescent growth spurt has long been recognized [7,8]. Mismatched rapid  
64 growth in the long bones relative to muscular lengthening may disrupt structure, neuromuscular  
65 function, and physical performance [9].

66 Deehan et al. [10], state that an increased participation in sports predisposes the immature  
67 skeleton to injury. Furthermore, participation in high intensity sport entails an inherent risk of  
68 sports-related injuries, and this is heightened at various stages of growth and maturation [11].  
69 Maturation induces profound changes in the skeletal, neuromuscular, and tendinous systems of  
70 young athletes [12] and mismatches in biological maturity may create competitive inequality and  
71 increase the risk of injury [13]. Le Gall et al. [14], further point out that injury rates generally  
72 increase with increasing CA. However, CA is a poor indicator of biological maturity [15];  
73 moreover, Ardern et al. [16], report that chronological age alone is an unreliable indicator of  
74 skeletal maturity. Skeletal age (SA) is generally accepted as the most accurate method of  
75 assessing biological maturity [6,17], by identifying critical periods of development; it also offers a  
76 rational method for monitored age-specific training. Before initiating any program for mitigating

1  
2  
3 77 sports injuries, the magnitude of the problem must be identified and the extent of the injury  
4  
5 78 defined in terms of incidence and severity [18].  
6  
7  
8 79 A number of studies have been conducted involving injuries in adolescent footballers;  
9  
10 80 conversely, few studies have focused on injuries in non-footballer adolescent athletes in high  
11  
12 81 performance sporting environments [19]. Studies of anthropometric characteristics and  
13  
14 82 biological maturity status as injury risk factors in Middle-Eastern youths are also limited,  
15  
16 83 highlighting the need for more research in this area. Therefore, the purpose of the present study  
17  
18 84 was to investigate injury incidence according to biological maturity in highly trained youth  
19  
20 85 athletes based at a Middle Eastern Sports Academy.  
21  
22

23 86 **METHODS**  
24  
25

26 87 Sixty-seven highly trained adolescent athletes (age range 11–18 y) representing athletics and  
27  
28 88 racquet sports (table tennis and squash) from a Middle Eastern sports school were included in  
29  
30 89 this four-year study. A prospective, longitudinal cohort design was used and included separate  
31  
32 90 observation periods over four consecutive seasons (20010–2011, 2011–2012, 2012–2013, and  
33  
34 91 2013–2014), i.e., school years, which lasted from the beginning of September until the end of  
35  
36 92 June (~40 weeks). Participant maturity assessments included both anthropometric  
37  
38 93 measurements, collected three times a season, and SA assessments using Fels method  
39  
40 94 completed once, at the start of every season. Medical screening was performed at the  
41  
42 95 beginning of each season to determine health and injury status. All selected athletes had  
43  
44 96 clearance from a physician to participate in their respective sport. Written informed consent was  
45  
46 97 sought and obtained from parents and assent from all participants. The study was part of a  
47  
48 98 general sports science provision to the sports academy, and all procedures were reviewed and  
49  
50 99 granted by the Institutional Review Board (IRB) for Human Subjects and conformed to the  
51  
52 100 recommendations of the Declaration of Helsinki.  
53  
54  
55  
56  
57  
58  
59  
60

## 101 **Participants involvement**

102 Figure 1 shows the flow of participants in the study over consecutive seasons. A total of four  
103 sporting disciplines were analysed, grouped into two categories: athletics and fencing and  
104 racquet sports (squash and table tennis). This classification was based on specific sport  
105 characteristics and injury risk [20,21]. Inclusion criteria were as follows: (1) the athlete had to be  
106 enrolled in the sports school during at least one full school year; (2) athletes with injuries in  
107 previous seasons were not excluded from this study, but injuries present at the beginning of the  
108 observation period were not included in statistical analyses; and (3) injuries that were not  
109 sustained in the context of the sports program or data related to sickness or other general  
110 medical conditions were not used for further analysis.

111 <<<Insert Figure 1 here>>>

## 113 **Injury data collection**

114 All injuries were assessed by a physical therapist (AR) with experience of working within youth  
115 sport. Data from medical records were used to document all sports related injuries during the  
116 study. Each sporting discipline had a dedicated full-time physiotherapist and a full-time  
117 employed medical doctor at the sports academy. The medical record used an injury reporting  
118 system based upon the football injury reporting system [22] and the Sport Medicine Diagnostic  
119 Coding System [23]. Information was gathered concerning all injuries related to sports activity,  
120 including several related variables (e.g. type, location, affected structure, mechanism [acute vs.  
121 overuse], time loss, severity, and date of injury).

## 122 **Somatic maturation and anthropometric measurements**

123 Anthropometric measurements were initially carried out on all participants on a three monthly  
124 basis along with an estimation of the age at peak height velocity (APHV) as a relative indicator



of somatic maturity and representing the time of maximum growth in stature during adolescence [1]. To ensure that the outcome measures remained consistent and reliable, every effort was made to ensure that measurements were taken at approximately the same time of the season. Measurements were collected by qualified practitioners from the International Society for the Advancement of Kinanthropometry (ISAK) and included stretch stature ( $\pm 0.1$  cm Holtain Limited, Crosswell, UK).

The predicted mature height (PMH) of all participants were collected and categorized into four PMH quartiles (Q1–Q4: Q1,  $<176$  cm; Q2, 176–180 cm; Q3, 180–184 cm; Q4,  $\geq 184$  cm). The athletes were then divided into three maturity groups (late, normal, or early maturing) based on the mean  $\pm 1.0$  year of the APHV of the total sample (late, APHV  $>$  mean + 1.0 y; normal, APHV within mean  $\pm 1.0$  y; early, APHV  $<$  mean – 1.0 y). Years from peak height velocity (maturity offset value: CA – maturity offset) was calculated by subtracting the CA at the date of injury from the age at estimated peak height velocity.

**Skeletal maturation assessment**

Each year consenting athletes had a radiograph of the left wrist and hand, a convenient area to examine, and a more accurate method for the assessment of SA [9], using the Fels method [6,24] which has an advantage over other methods [25]. Maturity status, defined by the difference between CA and SA was calculated and classified into four categories: late, normal, early, and mature athletes. Late referred to an SA that was younger than CA by more than 1.0 y, athletes with a normal pattern of maturity had an SA that was within 1.0 y of CA, early referred to an SA that was older than CA by more than 1.0 y, and the closure of growth plate determine skeletally mature athletes.

**Definition of injury**

Injuries were recorded as a physical complaint requiring the attention of medical staff, which occurred during sports training, strength and conditioning training or during competition. Injuries were divided into time-loss (TL) injuries and no time-loss (NTL) injuries. A clinical examination and/or treatment of an athlete which did not result in a full training session or competition being missed was described as a “medical attention” with NTL injury. A clinical examination and/or treatment of an athlete resulting in a training session or competition being missed the following day(s) was labelled as a TL injury [22]. A traumatic injury was defined as any injury resulting from a specific and identifiable mechanism, including contact and non-contact circumstances with acute onset. Overuse injuries were defined as injuries resulting from insidious onset without a recognisable mechanism. Injury severity was defined, based on days of absence from usual sport participation, as slight (1 d or less), minimal (2–3 d), mild (4–7 d), moderately serious (8–28 d), serious (>28 d up to 6 months) or long-term (>6 months) in accordance with [26].

## Statistical Analysis

Data were analysed using statistical software (Stata Corporation, College Station, Texas). Descriptive statistics were presented as frequencies and proportions (%), and incidence rates were expressed as the number of injuries/number of registered athletes. To examine the role of growth status and maturity with the onset of injuries, a univariate Cox regression survival analysis was performed after accounting for repeated visits of some athletes over the four seasons. Hazard ratios (HR) with 95% confidence intervals (CIs) were reported for each factor. Kaplan-Meier curves were plotted for SA groups and time to injury over a season. Where appropriate, 95% CIs are presented. The alpha level of significance was set at 5%.

## RESULTS

Throughout the study period, 67 athletes were enrolled. Table 1 presents the anthropometric characteristics of participants and their maturity status. From these participants, 43 (64%)

reported one or more injuries adding up to 212 injuries in total. Over the four seasons, the injury rate observed was 4.9 per registered athlete.

**Table 1.** Anthropometric characteristics (mean ± SD) of participants according to maturity status

	Late	Normal	Early
	(n = 4, 6.0%)	(n = 59, 88.1%)	(n = 4, 6.0%)
CA (years)	13.3 ± 1.1	12.3 ± 1.0	12.1 ± 0.5
Years from PHV	-2.4 ± 1.2	-1.6 ± 1.1	-0.1 ± 0.9
APHV (years)	15.8 ± 1.5	13.9 ± 0.5	12.2 ± 0.9
PMH (cm)	181.6 ± 17.1	179.4 ± 4.9	188.4 ± 3.5
% PMH (%)	85.0 ± 3.0	85.0 ± 4.0	90.0 ± 4.0
SA (years)	11.8 ± 0.5	12.8 ± 1.5	12.7 ± 1.8

CA: chronological age; APHV: age at peak height velocity; PMH: predicted mature height; SA: skeletal age; SD: standard deviation

**Skeletal age: maturity status distribution and injury risk**

Among all participants (n = 67), 4% were classified as late maturers, 33% as normal, 41% as early and 22% as skeletally mature. The overall injury free survival analysis of maturity status using SA assessment indicated that early maturing athletes had a two-fold higher risk of injury over a season compared to late maturing athletes (HR 2.04, 95% CI 1.15–3.61, P = 0.015; Figure 2). There was a trend that early maturing athletes had a greater risk of injury over a season compared to normal athletes (HR 1.62, 95% CI 0.99–2.65, P = 0.053), but this was only marginally significant. However, injury risk among late and fully mature athletes did not differ from normal maturers.

<<<<Insert Figure 2 Here >>>>

**Somatic maturation and anthropometric measurements: distribution and injury risk**

Using anthropometric measurements, among all participants (n = 67), 6.0% were classified as late maturing, 85.8% as normal, and 6.0% as early. Classification of participant maturity status (late, normal, and early) according to age at PHV (APHV) was not significantly associated with

194 overall injury incidence in this cohort of highly trained Middle-Eastern youth athletes. Older  
195 PHVs were marginally associated with higher injury risk, but this was not statistically significant  
196 (HR 1.11, 95% CI 0.99–1.23,  $P = 0.067$ ).

197 Both PMH (cm), and %PMH (%) were found to be associated with injury risk (HR 1.05, 95% CI  
198 1.01–1.08,  $P = 0.006$ , and HR 1.03, 95% CI 1.00–1.06,  $P = 0.026$ ), respectively. When  
199 compared to participants in the 1st quartile for PMH (<176), athletes in the 4th quartile ( $\geq 184$   
200 cm) had a two and half times greater risk of injury (HR 2.41, 95% CI 1.42–4.08,  $P = 0.001$ ) over  
201 a season.

202 No significant differences were observed in the injury risk between racquet sports ( $n = 30$ ) and  
203 individual sports athletes ( $n = 37$ ; HR 1.14, 95% CI, 0.86–1.52,  $P = 0.37$ ).

## 204 DISCUSSION

205 The present investigation was carried out to examine injury incidence according to maturity  
206 status. Biological maturity status and height gradient play a significant role in injury risk profiles  
207 of highly trained youth athletes. The results of the current study show that athletes maturing at a  
208 younger age are at significantly greater risk of injury, more than two-fold, compared to their later  
209 maturing counterparts. Taller athletes were also found to be significantly more at risk of injury.

210 There is limited and contrasting evidence on the relationship between maturity and injury in  
211 youth sports [27–29]. In this study, SA maturity (Fels method) showed that early maturing  
212 athletes had twice the risk of injury over a season compared to late maturing athletes. This  
213 finding is consistent with previous study [6], that described that early maturing athletes are  
214 significantly more at risk of injury than late or normally maturing athletes. A possible explanation  
215 could be that youth players with higher engagement and performance advantages are often  
216 associated with early maturation, usually transient during adolescence, and maybe reversed in  
217 early adulthood.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

218 However, our study results were inconsistent with other study [28] on youth athletes, in which  
219 late maturing athletes have a higher injury rate compared to their earlier maturing counterparts.  
220 A plausible explanation could be that Fourchet et al. [28], examined anthropometric data  
221 collected from a track and field cohort for their findings, while our study resulted from maturity  
222 status derived from bone age.

223 In the present study, no significant association was observed between APHV and injury risk  
224 (HR 0.90, 95% CI 0.74–1.11, P = 0.329), which is inconsistent with recent data on youth alpine  
225 ski racing [30] and other studies on talented Dutch and English youth soccer players [6,31]  
226 which show a heightened period of risk around the time of peak height velocity. An explanation  
227 of these discrepancies could be that our study cohort was not large enough, as the APHV  
228 method appears to be useful in youth talent selection and injury prevention programs because it  
229 can be easily applied in a large cohort of young athletes [32].

230 PMH and %PMH at a given age are minimally invasive, feasibly practical indicators of somatic  
231 maturation [15,33], especially if mature height can be assessed without an estimate of SA [24].  
232 In this study, the PMH and %PMH revealed that both indicators were associated with injury risk  
233 (HR 1.05, 95% CI 1.01–1.08, P = 0.006), and HR 1.03, 95% CI 1.00–1.06, P = 0.026),  
234 respectively. When compared to participants in the 1st quartile for PMH (<176), athletes in the  
235 4th quartile (≥184 cm) had two and a half times greater risk of injury (HR 2.41, 95% CI 1.42–  
236 4.08, P = 0.001). The present results are partly in line with previous studies on other sports.  
237 Johnson et al. [6], showed that gains in height in youth footballers over a season were  
238 associated with an increased number of injuries. The study of Kemper et al. [34], on elite youth  
239 soccer players with growth rates of at least 0.6 cm/month showed a higher risk for injury. In a  
240 different study on soccer athletes, it was found that the tallest boys had the highest incidence of  
241 injury [35]. However, these findings and those of the present study are not in line with a study on  
242 youth football players [36], in which injured and non-injured players did not differ in percentage

of mature height. An explanation could be that the definition of reportable injury in the methods of the study, which considered only time loss injuries, did not capture the full spectrum of injuries and therefore overlooked other injuries with insidious onset e.g. growth conditions.

The results of this study have some important practical implications. Malina et al. [2], advocate the documentation of anthropometric characteristics, biological maturity, and physical fitness parameters as crucial aids in the prevention of injury. Noninvasive methods for estimating maturity status may allow youth programs to match players using maturity status rather than CA, and thus equalize competition to some extent. An unequal competition is regarded as an impediment to personal development [37]. Furthermore, it has been suggested that there is an overwhelming bias in sport favoring taller athletes [38], and data on Olympic medal winners show that many running and jumping events are seriously biased in favor of the very tall [39].

When examining the classification resulting from SA of late (4%), normal (33%), early (41%), and skeletally mature athletes (22%), the under-representation of late and preponderance of early maturing athletes in this cohort is consistent with observations for male youth athletes in several sports including soccer and alpine ski racing [17,27,30]. However, these results and those of the present study are not in line with the study of Johnson et al. [6], on schoolboy footballers, in which two thirds of their players fall within the normal maturity category. Moreover, Le Gall et al. [14], classify only 12.0% as late maturers, 63.5% as normal maturers, and 24.5% as early maturers. These discrepancies are believed to be due to differences in selection policies and talent identification policies (physical, technical, and tactical skills) of varying elite development centers. Several studies point out that athletes who are more advanced in their biological maturity perform better than their later maturing peers and have a better chance of being selected [40–42]. Youth sport is highly selective, with a maturity-associated selection/exclusion process [33].

## Implications and concepts for prevention

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The findings in this study have several implications for youth athletes. First, our data suggests that adolescent athletes might be identified and selected with a preference for youths with advanced maturity. Such selection strategies which favor early maturers entail significant risks of injury. Accordingly, those involved in the selection and development of young athletes should be cognizant of temporary changes in motor control that may occur during these periods [43], consider maturity status, develop appropriate training programs to optimize training adaptation, design injury prevention plans to minimize activity related injury risk, and mitigate long term youth injury consequences.

Limitations of the current study should be noted. First, biological maturation methods have inherent limitations when applied to youth athletes and need to be applied with caution. Although SA is a gold standard indicator of maturation, it has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment protocols and the interpretation of results [44].

It must also be remembered that, except for accidents, a sports injury can rarely be ascribed to a single factor, but rather to an association of causes or circumstances and the interaction among a web of determinants [45,46].

**CONCLUSIONS**

The findings of the present study showed that maturity status plus PMH and %PMH are associated with injury in individual and racquet sports. As biological maturation varies individually in rate and timing, and mismatches in maturity may create competitive inequality and increase injury incidence, it is suggested that biological maturity should be considered during training to help prevent injury. Given the peculiarity of youth athletes it is important to optimize the planning of training activities to further improve the understanding of the link between training, growth, and injury.



## Figure legends

<<<Figure 1. Flowchart describing the inclusion and flow of participants throughout the study >>>

<<<Figure 2. Kaplan-Meier survival analysis of injuries in relation to different skeletal age (SA) maturity status >>>

## REFERENCES

- 1 Iuliano-Burns S, Mirwald RL, Bailey DA. Timing and magnitude of peak height velocity and peak tissue velocities for early, average, and late maturing boys and girls. *Am J Hum Biol* 2001;**13**:1–8. doi:10.1002/1520-6300(200101/02)13:1<1::AID-AJHB1000>3.0.CO;2-S
- 2 Malina RM, Cumming SP, Morano PJ, *et al.* Maturity status of youth football players: A noninvasive estimate. *Med Sci Sports Exerc* 2005;**37**:1044–52. doi:10.1249/01.mss.0000171622.45134.cb
- 3 Meeuwisse W. Assessing Causation in Sport Injury: A Multifactorial Model. *Clin J Sport Med* 1994;**4**:166–70. doi:10.1097/00042752-199407000-00004
- 4 Krivickas LS, Feinberg JH. Lower extremity injuries in college athletes: Relation between ligamentous laxity and lower extremity muscle tightness. *Arch Phys Med Rehabil* 1996;**77**:1139–43. doi:10.1016/S0003-9993(96)90137-9
- 5 Aicardi G, Vignolo M, Milani S, *et al.* Assessment of skeletal maturity of the hand-wrist and knee: A comparison among methods. *Am J Hum Biol* 2000;**12**:610–5. doi:10.1002/1520-6300(200009/10)12:5<610::AID-AJHB5>3.0.CO;2-D
- 6 Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. *Bmj* 2009;**338**:b490–b490. doi:10.1136/bmj.b490



1  
2  
3 316 7 Peterson HA. *Epiphyseal growth plate fractures*. 2007. doi:10.1007/978-3-540-33802-4  
4  
5  
6 317 8 Bailey DA, Wedge JH, McCulloch RG, *et al*. Epidemiology of fractures of the distal end of  
7  
8 318 the radius in children as associated with growth. *J Bone Jt SurgAm* 1989;**71**:1225–  
9  
10 319 31.<http://www.ncbi.nlm.nih.gov/pubmed/2777851>  
11  
12  
13 320 9 Lloyd RS, Oliver JL, Faigenbaum AD, *et al*. Chronological Age vs. Biological Maturation.  
14  
15 321 *J Strength Cond Res* 2014;**28**:1454–64. doi:10.1519/JSC.0000000000000391  
16  
17  
18 322 10 Deehan DJ, Bell K, McCaskie AW. Adolescent musculoskeletal injuries in a football  
19  
20 323 academy. *J Bone Jt Surg - Br Vol* 2007;**89-B**:5–8. doi:10.1302/0301-620X.89B1.18427  
21  
22  
23 324 11 Read P, Oliver JL, De Ste Croix MBA, *et al*. Injury Risk Factors in Male Youth Soccer  
24  
25 325 Players. *Strength Cond J* 2015;**37**:1–7. doi:10.1519/SSC.0000000000000171  
26  
27  
28 326 12 Huey K, Robinson MM, Wiesinger H-P, *et al*. Imbalances in the Development of Muscle  
29  
30 327 and Tendon as Risk Factor for Tendinopathies in Youth Athletes: A Review of Current  
31  
32 328 Evidence and Concepts of Prevention. 2017;**8**:1–18. doi:10.3389/fphys.2017.00987  
33  
34  
35 329 13 Cumming SP, Brown DJ, Mitchell S, *et al*. Premier League academy soccer players'  
36  
37 330 experiences of competing in a tournament bio-banded for biological maturation. *J. Sports*  
38  
39 331 *Sci*. 2017;;1–9. doi:10.1080/02640414.2017.1340656  
40  
41  
42 332 14 Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. *Scand J*  
43  
44 333 *Med Sci Sport* 2007;**17**:564–72. doi:10.1111/j.1600-0838.2006.00594.x  
45  
46  
47 334 15 Beunen GP, Rogol AD, Malina RM. Indicators of biological maturation and secular  
48  
49 335 changes in biological maturation. *Food Nutr. Bull*. 2006;**27**.  
50  
51 336 doi:10.1177/156482650903000307  
52  
53  
54 337 16 Ardern CL, Ekås G, Grindem H, *et al*. 2018 International Olympic Committee consensus  
55  
56 338 statement on prevention, diagnosis and management of paediatric anterior cruciate  
57  
58  
59  
60

- 339 ligament (ACL) injuries. *Knee Surgery, Sport Traumatol Arthrosc* 2018;**0**:1–17.
- 340 doi:10.1007/s00167-018-4865-y
- 341 17 Malina RM. Early sport specialization: Roots, effectiveness, risks. *Curr Sports Med Rep*
- 342 2010;**9**:364–71. doi:10.1249/JSR.0b013e3181fe3166
- 343 18 Leppänen M, Lapinleimu H, Lehtonen L, *et al*. Growth of extremely preterm infants born
- 344 in 2001-2010. *Acta Paediatr Int J Paediatr* 2013;**102**:206–8. doi:10.1111/apa.12061
- 345 19 Steffen K. More data needed on injury risk among young elite athletes. *Br J Sports Med*
- 346 2010;**44**:485–
- 347 9.<http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=51172712&site=ehost-live>
- 348 t-live
- 349 20 Rejeb A, Johnson A, Vaeyens R, *et al*. Compelling overuse injury incidence in youth
- 350 multisport athletes. *Eur J Sport Sci* 2017;**17**:495–502.
- 351 doi:10.1080/17461391.2016.1275820
- 352 21 Malisoux L, Frisch A, Urhausen A, *et al*. Injury incidence in a sports school during a 3-
- 353 year follow-up. *Knee Surgery, Sport Traumatol Arthrosc* 2013;**21**:2895–900.
- 354 doi:10.1007/s00167-012-2185-1
- 355 22 Fuller CW, Ekstrand J, Junge A, *et al*. Consensus statement on injury definitions and data
- 356 collection procedures in studies of football (soccer) injuries. *Scand. J. Med. Sci. Sport.*
- 357 2006;**16**:83–92. doi:10.1111/j.1600-0838.2006.00528.x
- 358 23 Meeuwisse W, H MD P, Tyreman H, *et al*. A Dynamic Model of Etiology in Sport Injury:
- 359 The Recursive Nature of Risk and Causation. *Clin J Sport Med* 2007;**17**:215–9.
- 360 doi:10.1097/JSM.0b013e3180592a48
- 361 24 Khamis HJ, Roche AF. Predicting adult stature without using skeletal age: the Khamis-

1  
2  
3 362 Roche method. *Pediatrics*. 1994;**94**:504–7. doi:10.1016/S0740-624X(98)90020-X  
4  
5  
6 363 25 Silventoinen K, Pke M, Tynelius P. Muscle Strength and Body Size and Later  
7  
8 364 Cerebrovascular and Coronary Heart Disease Maturity Status and Injury Risk in Youth  
9  
10 365 Soccer Players. *Sport Med* 2010;**20**:131–5. doi:10.1136/bmj.b490.Objective  
11  
12  
13 366 26 Timpka T, Alonso J-M, Jacobsson J, *et al*. Injury and illness definitions and data  
14  
15 367 collection procedures for use in epidemiological studies in Athletics (track and field):  
16  
17 368 Consensus statement. *Br J Sports Med* 2014;**48**:483–90. doi:10.1136/bjsports-2013-  
18  
19 369 093241  
20  
21  
22 370 27 Materne O, Farooq A, Johnson A, *et al*. Relationship between injuries and somatic  
23  
24 371 maturation in highly trained youth soccer players. *Int Res Sci Soccer II* 2016;;182–  
25  
26 372 92.[https://books.google.com/books?hl=en&lr=&id=w8DMCgAAQBAJ&oi=fnd&pg=PA182](https://books.google.com/books?hl=en&lr=&id=w8DMCgAAQBAJ&oi=fnd&pg=PA182&dq=Relationship+between+injuries+and+somatic+maturation+in+highly+trained+youth+soccer+players&ots=6D7Xy7rLzj&sig=ywUDNrV1u4pMrJIBN3wERj1u93M%5Cnhttp://s3.amazonaws.com/academia.edu)  
27  
28 373 &dq=Relationship+between+injuries+and+somatic+maturation+in+highly+trained+youth+  
29  
30 374 soccer+players&ots=6D7Xy7rLzj&sig=ywUDNrV1u4pMrJIBN3wERj1u93M%5Cnhttp://s3.  
31  
32 375 amazonaws.com/academia.edu.  
33  
34  
35 376 28 Fourchet F, Horobeanu C, Loepelt H, *et al*. Foot, Ankle, and Lower Leg Injuries in Young  
36  
37 377 Male Track and Field Athletes. *Int J Athl Ther Train* 2011;**16**:19–23 5p.  
38  
39 378 doi:10.1123/ijatt.16.3.19  
40  
41  
42 379 29 Swain M, Kamper SJ, Maher CG, *et al*. Relationship between growth, maturation and  
43  
44 380 musculoskeletal conditions in adolescents: a systematic review. *Br J Sports Med*  
45  
46 381 2018;;bjsports-2017-098418. doi:10.1136/bjsports-2017-098418  
47  
48  
49 382 30 Müller L, Hildebrandt C, Müller E, *et al*. Long-Term athletic development in youth alpine  
50  
51 383 ski racing: The effect of physical fitness, ski racing technique, anthropometrics and  
52  
53 384 biological maturity status on injuries. *Front Physiol* 2017;**8**. doi:10.3389/fphys.2017.00656  
54  
55  
56  
57  
58  
59  
60

- 385 31 Vanderlei FM, Vanderlei LCM, Bastos FN, *et al.* Characteristics and associated factors  
386 with sports injuries among children and adolescents. *Brazilian J Phys Ther* 2014;**18**:530–  
387 7. doi:10.1590/bjpt-rbf.2014.0059
- 388 32 Müller L. Müller EHCKK& RC. Die Erhebung des biologischen Entwicklungsstandes für  
389 die Talentselektion - welche Methode eignet sich? [The assessment of biological  
390 maturation for talent selection - which method can be used?]. *Sport Sport* 2015;**29**:56–63.
- 391 33 Bergeron MF, Mountjoy M, Armstrong N, *et al.* International Olympic Committee  
392 consensus statement on youth athletic development. *Br J Sports Med* 2015;**49**:843–51.  
393 doi:10.1136/bjsports-2015-094962
- 394 34 Kemper GLJ, Van Der Sluis A, Brink MS, *et al.* Anthropometric Injury Risk Factors in  
395 Elite-standard Youth Soccer. *Int J Sports Med* 2015;**36**:1112–7. doi:10.1055/s-0035-  
396 1555778
- 397 35 Backous DD, Friedl KE, Smith NJ, *et al.* Soccer Injuries and Their Relation to Physical  
398 Maturity. *Am J Dis Child* 1988;**142**:839–42. doi:10.1001/archpedi.1988.02150080045019
- 399 36 Malina RM, Morano PJ, Barron M, *et al.* Incidence and player risk factors for injury in  
400 youth football. *Clin J Sport Med* 2006;**16**:214–22. doi:10.1097/00042752-200605000-  
401 00005
- 402 37 Musch J, Grondin S. Unequal Competition as an Impediment to Personal Development: A  
403 Review of the Relative Age Effect in Sport. *Dev Rev* 2001;**21**:147–67.  
404 doi:10.1006/drev.2000.0516
- 405 38 Baxter-Jones ADG. Growth and Development of Young Athletes: Should Competition  
406 Levels be Age Related? *Sport Med* 1995;**20**:59–64. doi:10.2165/00007256-199520020-  
407 00001

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

408 39 Khosla T. Unfairness of certain events in the Olympic games. *Br Med J* 1968;**4**:111–3.

409 doi:10.1136/bmj.4.5623.111

410 40 Beunen GP, Malina RM, Van't Hof MA, *et al.* *Adolescent Growth and Motor Performance.*

411 1988.

412 41 Philippaerts RM, Vaeyens R, Janssens M, *et al.* The relationship between peak height

413 velocity and physical performance in youth soccer players. *J Sports Sci* 2006;**24**:221–30.

414 doi:10.1080/02640410500189371

415 42 Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity. *Growth,*

416 *Matur Phys Perform* 2004;:1–17.

417 43 Mueller L, Hildebrandt C, Mueller E, *et al.* Injuries and illnesses in a cohort of elite youth

418 alpine ski racers and the influence of biological maturity and relative age: a two-season

419 prospective study. *Open Access J Sport Med* 2017;**Volume 8**:113–22.

420 doi:10.2147/OAJSM.S133811

421 44 Malina RM, Rogol AD, Cumming SP, *et al.* Biological maturation of youth athletes:

422 assessment and implications. *Br J Sports Med* 2015;**49**:852–9. doi:10.1136/bjsports-

423 2015-094623

424 45 Theisen D, Malisoux L, Seil R, *et al.* : Injuries in Youth Sports: Epidemiology, Risk

425 Factors and Prevention. *Dtsch Z Sportmed* 2014;**65**:S248-252.[http://www.zeitschrift-](http://www.zeitschrift-sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-risk-factors-and-prevention/)

426 [sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-](http://www.zeitschrift-sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-risk-factors-and-prevention/)

427 [risk-factors-and-prevention/](http://www.zeitschrift-sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-risk-factors-and-prevention/)

428 46 Bittencourt NFN, Meeuwisse WH, Mendonça LD, *et al.* Complex systems approach for

429 sports injuries: moving from risk factor identification to injury pattern recognition—

430 narrative review and new concept. *Br J Sports Med* 2016;**50**:1309–14.

431 doi:10.1136/bjsports-2015-095850

432

433

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Contributors** AR and EW designed and developed the research question. AR, AJ and EW were involved in study design. AF supervised the data collection. AF analysed and interpreted the data. AR wrote the manuscript. All authors have contributed to and edited the manuscript and have approved the final manuscript.

**Funding** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Ethics approval** The study received ethical approval by the Anti-doping Lab Qatar

**Data sharing statement** No additional data are available

BMJ Open athletes  
included in the  
study in **2010/11**

3 athletes left the sports program at  
end season

6 athletes continued with the sports program

21 new athletes enrolled

27 athletes available in **2011/12**

7 athletes left the sports program at  
end season

20 athletes continued with the sports  
program

26 new athletes enrolled

46 athletes available in **2012/13**


12 athletes left the sports program  
at end season

36 athletes continued with the sports  
program

11 new athletes enrolled

47 athletes available in **2013/14**





93x65mm (300 x 300 DPI)

**STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies***

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	Page 1 and Page 2.
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Pages 2 and 3.
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Pages 4.
Objectives	3	State specific objectives, including any prespecified hypotheses	Pages 4 and 5.
Methods			
Study design	4	Present key elements of study design early in the paper	Page 5, line of methods 82
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 5.
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Pages 5 and 6. Figure 1.
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	On pages 6, 7 and 8. Definition and data collection of outcome variables were given.
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	On Pages 7 and 8.
Bias	9	Describe any efforts to address potential sources of bias	On Page 8.
Study size	10	Explain how the study size was arrived at	Every available athlete was included in our study.
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 5.

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Pages 8.
		(b) Describe any methods used to examine subgroups and interactions	Pages 5 and 6
		(c) Explain how missing data were addressed	Not applicable
		(d) If applicable, describe analytical methods taking account of sampling strategy	Not applicable
		(e) Describe any sensitivity analyses	Not applicable
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	All participants eligible completed the study. Page 7
		(b) Give reasons for non-participation at each stage	Not applicable
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	(Not applicable)
Outcome data	15*	Report numbers of outcome events or summary measures	Page 8 and page 9
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1.
		(b) Report category boundaries when continuous variables were categorized	Not applicable
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Not applicable
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Figure 2. different maturity level compared.
Discussion			
Key results	18	Summarise key results with reference to study objectives	Page 10.
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 13.
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 13.
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 13.
Other information			

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Not applicable
---------	----	---	----------------

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Sports Injuries Aligned to Predicted Mature Height in Highly Trained Middle-Eastern Youth Athletes: A Cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023284.R1
Article Type:	Research
Date Submitted by the Author:	25-Sep-2018
Complete List of Authors:	Rejeb, Abdallah; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie Johnson, Amanda; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center Farooq, Abdulaziz; Aspetar, Orthopaedic and Sports Medicine Department, Athlete Health and Performance Research Verrelest, Ruth ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie Pullinger, Samuel; ASPIRE Academy for Sports Excellence, Sports Sciences Vaeyens, Roel ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Department of movement and Sports Sciences Witvrouw, Erik; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapy
<b>Primary Subject Heading</b>:	Sports and exercise medicine
Secondary Subject Heading:	Paediatrics
Keywords:	Youth, Biological maturation, Skeletal age, Anthropometrics, Sports injury, Mature height

SCHOLARONE™  
Manuscripts

**Sports Injuries Aligned to Predicted Mature Height in Highly Trained Middle-Eastern Youth Athletes: A Cohort study**

**Authors:**

Abdallah Rejeb<sup>1 2</sup>, Amanda Johnson<sup>1</sup>, Abdulaziz Farooq<sup>1</sup>, Ruth Verrelest<sup>2</sup>, Samuel Pullinger<sup>3</sup>,  
Roel Vaeyens<sup>4</sup>, Erik Witvrouw<sup>2</sup>

**Affiliations:**

<sup>1</sup> Aspetar Orthopedic Sports Medicine and Hospital, Doha, Qatar

<sup>2</sup> Ghent University, Department of Rehabilitation Sciences and Physiotherapy, Belgium

<sup>3</sup> Aspire Academy for Sports Excellence, Doha, Qatar

<sup>4</sup> Ghent University, Department of Movement and Sports Sciences, Belgium

**Corresponding author**

Abdallah Rejeb

Address: Asian Games 2006 Road, P.O. Box 22287, Aspire Academy for Sports, Doha, Qatar

Phone Number: +974 66230638

Email: [Abdallah.rajeb@aspetar.com](mailto:Abdallah.rajeb@aspetar.com)

Manuscript word count = 3201

**ABSTRACT** (296/300 words)

**OBJECTIVES:** To investigate the association of maturity status with injury incidence in Middle-Eastern youth athletes.

**DESIGN:** Prospective cohort study

**SETTING:** Four consecutive seasons (2010 to 2014), Aspire Academy, Qatar.

**PARICIPANTS:** Male athletes (age range: 11–18 y) representing four disciplines enrolled and grouped into two categories: individual sports and racquet sports.

**OUTCOME MEASURES:** Injury data collected over four seasons. Athletes' anthropometric characteristics assessed to calculate APHV. Predicted mature heights (PMHs) collected and categorized into four quartiles. Athletes had wrist and hand radiographs for assessment of skeletal age (SA). Early and late maturers with an SA of >1 y older or younger than their chronological age (CA).

**RESULTS:** For the sample (n = 67) across all groups, 43 (64%) athletes had one or more injuries: total of 212 injuries, 4.9 injuries per athlete across study. Survival analysis of maturity status using SA found early maturing athletes had two-fold greater injury risk compared to late maturers ( [HR]; 2.04, 95% CI 1.15–3.61, P = 0.015). PMH associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006).

Athletes in 4th quartile ( $\geq 184$  cm) had up to 2-fold injury risk (HR 2.41, 95% CI 1.42–4.08, P = 0.001). Racquet and individual sports involved similar injury risk (HR 1.14, 95% CI 0.86–1.52, P = 0.37).

**CONCLUSION:** SA early maturity and PMH gradient were significant predictors of injury in youths.

**STRENGTHS AND LIMITATIONS OF THE STUDY**

- First longitudinal study to assess anthropometric characteristics and biological maturity status as injury risk factors in Middle-Eastern athletes.
- Participants were highly trained adolescent athletes.
- Measurement of maturity and growth were at moderate-to-high risk of bias.
- SA has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment and interpretation of results.

**INTRODUCTION**

The range of somatic and biological maturity in individuals of the same chronological age (CA) is large [1]. Such observations are derived from correlational and multivariate studies that compare young individuals of the same age who are at both extremes of the maturity range [2]. Therefore, the assessment of maturity is an important consideration when dealing with adolescent athletes on a longitudinal basis. Further, understanding the cause of disease and injury is vital in predicting and preventing injury [3].

In young athletes, the demands of their chosen sport are superimposed on normal growth and maturation. A literature review revealed that there is a greater susceptibility to injury during certain periods of growth [4–6]. Indeed, the association between an increased prevalence of injuries and the adolescent growth spurt has long been recognized [7–9]. A recent study analysis [10] on adolescent soccer players revealed greater risk of injury with players within age at peak height velocity (APHV) in comparison with the players before and after APHV. Mismatched rapid growth in the long bones relative to muscular lengthening may disrupt structure, neuromuscular function, and physical performance [11].

Deehan et al. [12], state that an increased participation in sports predisposes the immature skeleton to injury. Furthermore, participation in high intensity sport entails an inherent risk of



sports-related injuries, and this is heightened at various stages of growth and maturation [13]. Maturation induces profound changes in the skeletal, neuromuscular, and tendinous systems of young athletes [14] and mismatches in biological maturity may create competitive inequality and increase the risk of injury [15]. Le Gall et al. [16], further point out that injury rates generally increase with increasing CA. However, CA is a poor indicator of biological maturity [17]; moreover, Ardern et al. [18], report that chronological age alone is an unreliable indicator of skeletal maturity. Skeletal age (SA) is generally accepted as the most accurate method of assessing biological maturity [6,19], by identifying critical periods of development; it also offers a rational method for monitored age-specific training. Before initiating any program for mitigating sports injuries, the magnitude of the problem must be identified and the extent of the injury defined in terms of incidence and severity [20].

A number of studies have been conducted involving injuries in adolescent footballers; conversely, few studies have focused on injuries in non-footballer adolescent athletes in high performance sporting environments [21]. Studies of anthropometric characteristics and biological maturity status as injury risk factors in Middle-Eastern youths are also limited, highlighting the need for more research in this area. Therefore, the purpose of the present study was to investigate injury incidence according to biological maturity using two outcome measures (SA and PHV) in highly trained youth athletes based at a Middle Eastern Sports Academy.

## METHODS

Sixty-seven highly trained adolescent athletes (age range 11–18 y) representing athletics and racquet sports (table tennis and squash) from a Middle Eastern sports school were included in this four-year study. A prospective, longitudinal cohort design was used and included separate observation periods over four consecutive seasons (2010–2011, 2011–2012, 2012–2013, and 2013–2014), i.e., school years, which lasted from the beginning of September until the end of June (~40 weeks). Participant maturity assessments included both anthropometric

measurements, collected three times a season, and SA assessments using Fels method completed once, at the start of every season. Medical screening was performed at the beginning of each season to determine health and injury status. All selected athletes had clearance from a physician to participate in their respective sport. Written informed consent was sought and obtained from parents and assent from all participants. The study was part of a general sports science provision to the sports academy, and all procedures were reviewed and granted by the Institutional Review Board (IRB) for Human Subjects and conformed to the recommendations of the Declaration of Helsinki.

**Participants**

Figure 1 shows the flow of participants in the study over consecutive seasons. A total of four sporting disciplines were analysed, grouped into two categories: athletics and fencing and racquet sports (squash and table tennis). This classification was based on specific sport characteristics and injury risk [22,23]. Inclusion criteria were as follows: (1) the athlete had to be enrolled in the sports school during at least one full school year; (2) athletes with injuries in previous seasons were not excluded from this study, but injuries present at the beginning of the observation period were not included in statistical analyses; and (3) injuries that were not sustained in the context of the sports program (e.g. recreational activities) or data related to sickness or other general medical conditions were not used for further analysis.

<<<Insert Figure 1 here>>>

**Injury definition and data collection**

An Injury was defined as any physical complaint, which occurred during sports training, strength and conditioning training or during competition. Injuries were divided into time-loss (TL) injuries and no time-loss (NTL) injuries. A clinical examination and/or treatment of an athlete which did

not result in a full training session or competition being missed was described as a complaint with NTL injury. A clinical examination and/or treatment of an athlete resulting in a training session or competition being missed the following day(s) was labelled as a TL injury [23]. A traumatic injury was defined as any injury resulting from a specific and identifiable mechanism, including contact and non-contact circumstances with acute onset. Overuse injuries were defined as injuries resulting from insidious onset without a recognizable mechanism. Injury severity was defined, based on days of absence from usual sport participation, as slight (1 d or less), minimal (2–3 d), mild (4–7 d), moderately serious (8–28 d), serious (>28 d up to 6 months) or long-term (>6 months) in accordance with [24].

All injuries were collected by a physical therapist (AR) with experience of working within youth sport. Data from medical records were used to document all sports related injuries during the study. Each sporting discipline had a dedicated full-time physiotherapist and a full-time employed medical doctor at the sports academy. The medical record used an injury reporting system based upon the football injury reporting system [25] and the Sport Medicine Diagnostic Coding System [26]. Information was gathered concerning all injuries related to sports activity, including several related variables (e.g. type, location, affected structure, mechanism [acute vs. overuse], time loss, severity, and date of injury).

### **Somatic maturation and anthropometric measurements**

Anthropometric measurements were initially carried out on all participants on a three monthly basis along with an estimation of the age at peak height velocity (APHV) as a relative indicator of somatic maturity and representing the time of maximum growth in stature during adolescence using Mirwald method [27] for the prediction of growth [1]. APHV was calculated from the first measurement recorded. To ensure that the outcome measures remained consistent and reliable, every effort was made to ensure that measurements were taken at approximately the same time of the season. Measurements were collected by qualified practitioners from the

International Society for the Advancement of Kinanthropometry (ISAK) and included stretch stature ( $\pm 0.1$  cm Holtain Limited, Crosswell, UK).

The predicted mature height (PMH) of all participants were collected and categorized into four PMH quartiles (Q1–Q4: Q1 < 176 cm; 176 cm  $\leq$  Q2 < 180 cm; 180 cm  $\leq$  Q3 < 184 cm; Q4  $\geq$  184 cm). The athletes were then divided into three maturity groups (late, normal, or early maturing) based on the mean  $\pm 1.0$  year of the APHV of the total sample (late, APHV > mean + 1.0 y; normal, APHV within mean  $\pm 1.0$  y; early, APHV < mean – 1.0 y). Years from peak height velocity (maturity offset value: CA – maturity offset) was calculated by subtracting the CA at the date of injury from the age at estimated peak height velocity.

**Skeletal maturation assessment**

Each year athletes were required to have a radiograph of the left wrist and hand, a convenient area to examine, and a more accurate method for the assessment of SA [11], using the Fels method [6,28] which has an advantage over other methods [29]. Maturity status, defined by the difference between CA and SA was calculated and classified into four categories: late, normal, early, and mature athletes. Late referred to an SA that was younger than CA by more than 1.0 y, athletes with a normal pattern of maturity had an SA that was within 1.0 y of CA, early referred to an SA that was older than CA by more than 1.0 y, and the closure of growth plate determine skeletally mature athletes.

**Statistical Analysis**

Data were analysed using statistical software (Stata Corporation, College Station, Texas). Descriptive statistics were presented as frequencies and proportions (%), and incidence rates were expressed as the number of injuries/number of registered athletes. To examine the role of growth status and maturity with the onset of injuries, a univariate Cox regression survival analysis was performed after accounting for repeated visits of athletes over the four seasons.

Hazard ratios (HR) with 95% confidence intervals (CIs) were reported for each factor. Kaplan-Meier curves were plotted for SA groups and time to injury over a season. Where appropriate, 95% CIs are presented. The alpha level of significance was set at 5%.

### Patient and public involvement statement

Patients and public were not involved in the analysis of this study.

## RESULTS

Throughout the four-year seasons study period, 67 athletes were enrolled representing 151 athletic seasons. Table 1 presents the anthropometric characteristics of participants and their maturity status. From these participants, 43 (64%) reported one or more injuries adding up to 212 injuries in total. The injury rate observed per registered athlete amounted to 4.9 injuries over the course of four seasons.

**Table 1.** Anthropometric characteristics (mean  $\pm$  SD) of participants according to maturity status

	Late	Normal	Early
	(n = 4, 6.0%)	(n = 59, 88.1%)	(n = 4, 6.0%)
CA (years)	13.3 $\pm$ 1.1	12.3 $\pm$ 1.0	12.1 $\pm$ 0.5
Years from PHV	-2.4 $\pm$ 1.2	-1.6 $\pm$ 1.1	-0.1 $\pm$ 0.9
APHV (years)	15.8 $\pm$ 1.5	13.9 $\pm$ 0.5	12.2 $\pm$ 0.9
PMH (cm)	181.6 $\pm$ 17.1	179.4 $\pm$ 4.9	188.4 $\pm$ 3.5
% PMH (%)	85.0 $\pm$ 3.0	85.0 $\pm$ 4.0	90.0 $\pm$ 4.0
SA (years)	11.8 $\pm$ 0.5	12.8 $\pm$ 1.5	12.7 $\pm$ 1.8

CA: chronological age; APHV: age at peak height velocity; PMH: predicted mature height; SA: skeletal age; SD: standard deviation

### Skeletal age: maturity status distribution and injury risk

Among all participants (n = 67), 4% were classified as late maturers, 33% as normal, 41% as early and 22% as skeletally mature. The overall injury free survival analysis of maturity status using SA assessment indicated that early maturing athletes had a two-fold higher risk of injury over a season compared to late maturing athletes (HR 2.04, 95% CI 1.15–3.61, P = 0.015),

(Figure 2). There was a trend that early maturing athletes had a greater risk of injury over a season compared to normal athletes (HR 1.62, 95% CI 0.99–2.65, P = 0.053), but this was only marginally significant. However, injury risk among late and fully mature athletes did not differ from normal maturers.

<<<<Insert Figure 2 Here >>>>

**Somatic maturation and anthropometric measurements: distribution and injury risk**

Using anthropometric measurements, among all participants (n = 67), 6.0% were classified as late maturing, 85.8% as normal, and 6.0% as early. Classification of participant maturity status (late, normal, and early) according to age at PHV (APHV) was not significantly associated with overall injury incidence in this cohort of highly trained Middle-Eastern youth athletes. Older PHVs were marginally associated with higher injury risk, but this was not statistically significant (HR 1.11, 95% CI 0.99–1.23, P = 0.067).

Both PMH (cm), and %PMH were found to be associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006, and HR 1.03, 95% CI 1.00–1.06, P = 0.026), respectively. When compared to participants in the 1st quartile for PMH (<176), athletes in the 4th quartile (≥184 cm) had a two and half times greater risk of injury (HR 2.41, 95% CI 1.42–4.08, P = 0.001) over a season.

No significant differences were observed in the injury risk between racquet sports (n = 30) and individual sports athletes (n = 37; HR 1.14, 95% CI, 0.86–1.52, P = 0.37).

**DISCUSSION**

The present investigation was carried out to examine injury incidence according to maturity status. Biological maturity status and height gradient play a significant role in injury risk profiles of highly trained youth athletes. The results of the current study show that athletes maturing at a

1  
2  
3 younger age are at significantly greater risk of injury, more than two-fold, compared to their later  
4 maturing counterparts. Taller athletes were also found to be significantly more at risk of injury.  
5  
6

7  
8 There is limited and contrasting evidence on the relationship between maturity and injury in  
9 youth sports [10,30,31]. In this study, SA maturity (Fels method) showed that early maturing  
10 athletes had twice the risk of injury over a season compared to late maturing athletes. This  
11 finding is consistent with previous study [6], that described that early maturing athletes are  
12 significantly more at risk of injury than late or normally maturing athletes. A possible explanation  
13 could be that youth players with higher engagement and performance advantages are often  
14 associated with early maturation, usually transient during adolescence, and maybe reversed in  
15 early adulthood [16]  
16  
17

18  
19 However, our study results were inconsistent with other study [30] on youth athletes, in which  
20 late maturing athletes have a higher injury rate compared to their earlier maturing counterparts.  
21 A plausible explanation could be that Fourchet et al. [30], examined anthropometric data  
22 collected from a track and field cohort for their findings, while our study resulted from maturity  
23 status derived from bone age but with no substantial association from APHV.  
24  
25

26  
27 In the present study, no significant association was observed between APHV and injury risk  
28 (HR 0.90, 95% CI 0.74–1.11,  $P = 0.329$ ), which is inconsistent with recent data on youth alpine  
29 ski racing [32] and other studies on talented Dutch and English youth soccer players [6,33]  
30 which show a heightened period of risk around the time of peak height velocity. An explanation  
31 of these discrepancies could be that our study cohort was not large enough, as the APHV  
32 method appears to be useful in youth talent selection and injury prevention programs because it  
33 can be easily applied in a large cohort of young athletes [34].  
34  
35

36  
37 PMH and %PMH at a given age are minimally invasive, feasibly practical indicators of somatic  
38 maturation [17,35], especially if mature height can be assessed without an estimate of SA [28].  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

In this study, the PMH and %PMH revealed that both indicators were associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006), and HR 1.03, 95% CI 1.00–1.06, P = 0.026), respectively. When compared to participants in the 1st quartile for PMH (<176), athletes in the 4th quartile ( $\geq 184$  cm) had two and a half times greater risk of injury (HR 2.41, 95% CI 1.42–4.08, P = 0.001). The present results are partly in line with previous studies on other sports. Johnson et al. [6], showed that gains in height in youth footballers over a season were associated with an increased number of injuries. The study of Kemper et al. [36], on elite youth soccer players with growth rates of at least 0.6 cm/month showed a higher risk for injury. In a different study on soccer athletes, it was found that the tallest boys had the highest incidence of injury [37]. However, these findings and those of the present study are not in line with a study on youth football players [38], in which injured and non-injured players did not differ in percentage of mature height. An explanation could be that the definition of reportable injury in the methods of the study, which considered only time loss injuries, did not capture the full spectrum of injuries and therefore overlooked other injuries with insidious onset e.g. growth conditions.

The results of this study have some important practical implications. Malina et al. [2], advocate the documentation of anthropometric characteristics, biological maturity, and physical fitness parameters as crucial aids in the prevention of injury. Noninvasive methods for estimating maturity status may allow youth programs to match players using maturity status rather than CA, and thus equalize competition to some extent. An unequal competition is regarded as an impediment to personal development [39]. Furthermore, it has been suggested that there is an overwhelming bias in sport favoring taller athletes [40], and data on Olympic medal winners show that many running and jumping events are seriously biased in favor of the very tall [41].

When examining the classification resulting from SA of late (4%), normal (33%), early (41%), and skeletally mature athletes (22%), the under-representation of late and preponderance of early maturing athletes in this cohort is consistent with observations for male youth athletes in



several sports including soccer and alpine ski racing [10,19,32]. However, these results and those of the present study are not in line with the study of Johnson et al. [6], on schoolboy footballers, in which two thirds of their players fall within the normal maturity category. Moreover, Le Gall et al. [16], classify only 12.0% as late maturers, 63.5% as normal maturers, and 24.5% as early maturers. These discrepancies are believed to be due to differences in selection policies and talent identification policies (physical, technical, and tactical skills) of varying elite development centers. Several studies point out that athletes who are more advanced in their biological maturity perform better than their later maturing peers and have a better chance of being selected [42–44]. Youth sport is highly selective, with a maturity-associated selection/exclusion process [35].

### **Implications and concepts for prevention**

The findings in this study have several implications for youth athletes. First, our data suggests that adolescent athletes might be identified and selected with a preference for youths with advanced maturity. Such selection strategies which favor early maturers entail significant risks of injury. Accordingly, those involved in the selection and development of young athletes should be cognizant of temporary changes in motor control that may occur during these periods [45], consider maturity status, develop appropriate training programs to optimize training adaptation, design injury prevention plans to minimize activity related injury risk, and mitigate long term youth injury consequences.

Limitations of the current study should be noted. First, biological maturation methods have inherent limitations when applied to youth athletes and need to be applied with caution. Although SA is a gold standard indicator of maturation, it has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment protocols and the interpretation of results [46]. Although our sample size is small, we have a follow-up over four

seasons. Another limitation, we had no data on training or competition exposure, which reduces the comparability with other studies reporting injury incidence.

It must also be remembered that, except for accidents, a sports injury can rarely be ascribed to a single factor, but rather to an association of causes or circumstances and the interaction among a web of determinants [47,48].

CONCLUSIONS

The findings of the present study showed that maturity status plus PMH and %PMH are associated with injury in individual and racquet sports but no association has been established between APHV and injury. As SA varies individually in rate and timing, and mismatches in maturity may create competitive inequality and increase injury incidence, it is suggested that biological maturity should be considered during training to help prevent injury. Given the peculiarity of youth athletes it is important to optimize the planning of training activities to further improve the understanding of the link between training, growth, and injury.

Figure legends

<<<Figure 1. Flowchart describing the inclusion and flow of participants throughout the study >>>  
<<<Figure 2. Kaplan-Meier survival analysis of injuries in relation to different skeletal age (SA) maturity status >>>

REFERENCES

1 Iuliano-Burns S, Mirwald RL, Bailey DA. Timing and magnitude of peak height velocity and peak tissue velocities for early, average, and late maturing boys and girls. *Am J Hum Biol* 2001;**13**:1–8. doi:10.1002/1520-6300(200101/02)13:1<1::AID-AJHB1000>3.0.CO;2-S

2 Malina RM, Cumming SP, Morano PJ, et al. Maturity status of youth football players: A

- noninvasive estimate. *Med Sci Sports Exerc* 2005;**37**:1044–52.  
doi:10.1249/01.mss.0000171622.45134.cb
- 3 Meeuwisse W. Assessing Causation in Sport Injury: A Multifactorial Model. *Clin J Sport Med* 1994;**4**:166–70. doi:10.1097/00042752-199407000-00004
- 4 Krivickas LS, Feinberg JH. Lower extremity injuries in college athletes: Relation between ligamentous laxity and lower extremity muscle tightness. *Arch Phys Med Rehabil* 1996;**77**:1139–43. doi:10.1016/S0003-9993(96)90137-9
- 5 Aicardi G, Vignolo M, Milani S, *et al.* Assessment of skeletal maturity of the hand-wrist and knee: A comparison among methods. *Am J Hum Biol* 2000;**12**:610–5.  
doi:10.1002/1520-6300(200009/10)12:5<610::AID-AJHB5>3.0.CO;2-D
- 6 Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. *Bmj* 2009;**338**:b490–b490. doi:10.1136/bmj.b490
- 7 Peterson HA. *Epiphyseal growth plate fractures*. 2007. doi:10.1007/978-3-540-33802-4
- 8 Bailey DA, Wedge JH, McCulloch RG, *et al.* Epidemiology of fractures of the distal end of the radius in children as associated with growth. *J Bone Jt SurgAm* 1989;**71**:1225–31.  
<http://www.ncbi.nlm.nih.gov/pubmed/2777851>
- 9 Leppänen M, Pasanen K, Clarsen B, *et al.* Overuse injuries are prevalent in children's competitive football: a prospective study using the OSTRC Overuse Injury Questionnaire. *Br J Sports Med* 2018;**52**:bjsports-2018-099218. doi:10.1136/bjsports-2018-099218
- 10 Materne O, Farooq A, Johnson A, *et al.* Relationship between injuries and somatic maturation in highly trained youth soccer players. *Int Res Sci Soccer II* 2016;**1**:182–92.  
<https://books.google.com/books?hl=en&lr=&id=w8DMCgAAQBAJ&oi=fnd&pg=PA182>

&dq=Relationship+between+injuries+and+somatic+maturation+in+highly+trained+youth+soccer+players&ots=6D7Xy7rLzj&sig=ywUDNrV1u4pMrJIBN3wERj1u93M%5Cnhttp://s3.amazonaws.com/academia.edu.

11 Lloyd RS, Oliver JL, Faigenbaum AD, *et al.* Chronological Age vs. Biological Maturation. *J Strength Cond Res* 2014;**28**:1454–64. doi:10.1519/JSC.0000000000000391

12 Deehan DJ, Bell K, McCaskie AW. Adolescent musculoskeletal injuries in a football academy. *J Bone Jt Surg - Br Vol* 2007;**89-B**:5–8. doi:10.1302/0301-620X.89B1.18427

13 Read P, Oliver JL, De Ste Croix MBA, *et al.* Injury Risk Factors in Male Youth Soccer Players. *Strength Cond J* 2015;**37**:1–7. doi:10.1519/SSC.0000000000000171

14 Huey K, Robinson MM, Wiesinger H-P, *et al.* Imbalances in the Development of Muscle and Tendon as Risk Factor for Tendinopathies in Youth Athletes: A Review of Current Evidence and Concepts of Prevention. 2017;**8**:1–18. doi:10.3389/fphys.2017.00987

15 Cumming SP, Brown DJ, Mitchell S, *et al.* Premier League academy soccer players' experiences of competing in a tournament bio-banded for biological maturation. *J. Sports Sci.* 2017;;1–9. doi:10.1080/02640414.2017.1340656

16 Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. *Scand J Med Sci Sport* 2007;**17**:564–72. doi:10.1111/j.1600-0838.2006.00594.x

17 Beunen GP, Rogol AD, Malina RM. Indicators of biological maturation and secular changes in biological maturation. *Food Nutr. Bull.* 2006;**27**. doi:10.1177/156482650903000307

18 Ardern CL, Ekås G, Grindem H, *et al.* 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Knee Surgery, Sport Traumatol Arthrosc* 2018;**0**:1–17.

- doi:10.1007/s00167-018-4865-y
- 19 Malina RM. Early sport specialization: Roots, effectiveness, risks. *Curr Sports Med Rep* 2010;**9**:364–71. doi:10.1249/JSR.0b013e3181fe3166
- 20 Leppänen M, Lapinleimu H, Lehtonen L, *et al*. Growth of extremely preterm infants born in 2001-2010. *Acta Paediatr Int J Paediatr* 2013;**102**:206–8. doi:10.1111/apa.12061
- 21 Steffen K. More data needed on injury risk among young elite athletes. *Br J Sports Med* 2010;**44**:485–  
9.<http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=51172712&site=ehost-live>
- 22 Rejeb A, Johnson A, Vaeyens R, *et al*. Compelling overuse injury incidence in youth multisport athletes. *Eur J Sport Sci* 2017;**17**:495–502.  
doi:10.1080/17461391.2016.1275820
- 23 Malisoux L, Frisch A, Urhausen A, *et al*. Injury incidence in a sports school during a 3-year follow-up. *Knee Surgery, Sport Traumatol Arthrosc* 2013;**21**:2895–900.  
doi:10.1007/s00167-012-2185-1
- 24 Timpka T, Alonso J-M, Jacobsson J, *et al*. Injury and illness definitions and data collection procedures for use in epidemiological studies in Athletics (track and field): Consensus statement. *Br J Sports Med* 2014;**48**:483–90. doi:10.1136/bjsports-2013-093241
- 25 Fuller CW, Ekstrand J, Junge A, *et al*. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Scand. J. Med. Sci. Sport.* 2006;**16**:83–92. doi:10.1111/j.1600-0838.2006.00528.x
- 26 Meeuwisse W, H MD P, Tyreman H, *et al*. A Dynamic Model of Etiology in Sport Injury:

The Recursive Nature of Risk and Causation. *Clin J Sport Med* 2007;**17**:215–9.  
doi:10.1097/JSM.0b013e3180592a48

27 MIRWALD RL, G. BAXTER-JONES AD, BAILEY DA, *et al*. An assessment of maturity from anthropometric measurements. *Med Sci Sport Exerc* 2002;**34**:689–94.  
doi:10.1097/00005768-200204000-00020

28 Khamis HJ, Roche AF. Predicting adult stature without using skeletal age: the Khamis-Roche method. *Pediatrics*. 1994;**94**:504–7. doi:10.1016/S0740-624X(98)90020-X

29 Silventoinen K, Pke M, Tynelius P. Muscle Strength and Body Size and Later Cerebrovascular and Coronary Heart Disease Maturity Status and Injury Risk in Youth Soccer Players. *Sport Med* 2010;**20**:131–5. doi:10.1136/bmj.b490.Objective

30 Fourchet F, Horobeanu C, Loepelt H, *et al*. Foot, Ankle, and Lower Leg Injuries in Young Male Track and Field Athletes. *Int J Athl Ther Train* 2011;**16**:19–23 5p.  
doi:10.1123/ijatt.16.3.19

31 Swain M, Kamper SJ, Maher CG, *et al*. Relationship between growth, maturation and musculoskeletal conditions in adolescents: a systematic review. *Br J Sports Med* 2018;:bjssports-2017-098418. doi:10.1136/bjssports-2017-098418

32 Müller L, Hildebrandt C, Müller E, *et al*. Long-Term athletic development in youth alpine ski racing: The effect of physical fitness, ski racing technique, anthropometrics and biological maturity status on injuries. *Front Physiol* 2017;**8**. doi:10.3389/fphys.2017.00656

33 Vanderlei FM, Vanderlei LCM, Bastos FN, *et al*. Characteristics and associated factors with sports injuries among children and adolescents. *Brazilian J Phys Ther* 2014;**18**:530–7. doi:10.1590/bjpt-rbf.2014.0059

34 Müller L. Müller EHCKK& RC. Die Erhebung des biologischen Entwicklungsstandes für

- die Talentselektion - welche Methode eignet sich? [The assessment of biological maturation for talent selection - which method can be used?]. *Sport Sport* 2015;**29**:56–63.
- 35 Bergeron MF, Mountjoy M, Armstrong N, *et al.* International Olympic Committee consensus statement on youth athletic development. *Br J Sports Med* 2015;**49**:843–51. doi:10.1136/bjsports-2015-094962
- 36 Kemper GLJ, Van Der Sluis A, Brink MS, *et al.* Anthropometric Injury Risk Factors in Elite-standard Youth Soccer. *Int J Sports Med* 2015;**36**:1112–7. doi:10.1055/s-0035-1555778
- 37 Backous DD, Friedl KE, Smith NJ, *et al.* Soccer Injuries and Their Relation to Physical Maturity. *Am J Dis Child* 1988;**142**:839–42. doi:10.1001/archpedi.1988.02150080045019
- 38 Malina RM, Morano PJ, Barron M, *et al.* Incidence and player risk factors for injury in youth football. *Clin J Sport Med* 2006;**16**:214–22. doi:10.1097/00042752-200605000-00005
- 39 Musch J, Grondin S. Unequal Competition as an Impediment to Personal Development: A Review of the Relative Age Effect in Sport. *Dev Rev* 2001;**21**:147–67. doi:10.1006/drev.2000.0516
- 40 Baxter-Jones ADG. Growth and Development of Young Athletes: Should Competition Levels be Age Related? *Sport Med* 1995;**20**:59–64. doi:10.2165/00007256-199520020-00001
- 41 Khosla T. Unfairness of certain events in the Olympic games. *Br Med J* 1968;**4**:111–3. doi:10.1136/bmj.4.5623.111
- 42 Beunen GP, Malina RM, Van't Hof MA, *et al.* *Adolescent Growth and Motor Performance*. 1988.

43 Philippaerts RM, Vaeyens R, Janssens M, *et al.* The relationship between peak height velocity and physical performance in youth soccer players. *J Sports Sci* 2006;**24**:221–30. doi:10.1080/02640410500189371

44 Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity. *Growth, Matur Phys Perform* 2004;:1–17.

45 Mueller L, Hildebrandt C, Mueller E, *et al.* Injuries and illnesses in a cohort of elite youth alpine ski racers and the influence of biological maturity and relative age: a two-season prospective study. *Open Access J Sport Med* 2017;**Volume 8**:113–22. doi:10.2147/OAJSM.S133811

46 Malina RM, Rogol AD, Cumming SP, *et al.* Biological maturation of youth athletes: assessment and implications. *Br J Sports Med* 2015;**49**:852–9. doi:10.1136/bjsports-2015-094623

47 Theisen D, Malisoux L, Seil R, *et al.* : Injuries in Youth Sports: Epidemiology, Risk Factors and Prevention. *Dtsch Z Sportmed* 2014;**65**:S248-252. <http://www.zeitschrift-sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-risk-factors-and-prevention/>

48 Bittencourt NFN, Meeuwisse WH, Mendonça LD, *et al.* Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition—narrative review and new concept. *Br J Sports Med* 2016;**50**:1309–14. doi:10.1136/bjsports-2015-095850



**CONTRIBUTORSHIP STATEMENT:** AR designed and developed the research question and wrote the original version of the manuscript as part of his doctoral thesis. EW (doctoral supervisor) reviewed, designed and provided expertise to the study. AJ (doctoral committee member) was involved in study design. AF supervised and provided expertise with respect to the data analyses. VR, SP, and RV (doctoral committee member) reviewed and provided expertise to the study. All authors have contributed to and edited the manuscript and have approved the final manuscript.

**FUNDING:** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

**COMPETING INTERESTS:** None declared.

**Ethics approval:** The study received ethical approval by the Anti- Doping Lab Qatar

**DATA SHARING STATEMENT:** No additional data are available

---




Figure 1. Flowchart describing the inclusion and flow of participants throughout the study

135x89mm (300 x 300 DPI)




Figure 2. Kaplan-Meier survival analysis of injuries in relation to different skeletal age (SA) maturity status

96x67mm (300 x 300 DPI)

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	Page 1 and Page 2.
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Pages 2 and 3.
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Pages 4.
Objectives	3	State specific objectives, including any prespecified hypotheses	Pages 4 and 5.
Methods			
Study design	4	Present key elements of study design early in the paper	Page 5, line of methods 82
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 5.
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Pages 5 and 6. Figure 1.
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	On pages 6, 7 and 8. Definition and data collection of outcome variables were given.
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	On Pages 7 and 8.
Bias	9	Describe any efforts to address potential sources of bias	On Page 8.
Study size	10	Explain how the study size was arrived at	Every available athlete was included in our study.
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 5.

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Pages 8.
		(b) Describe any methods used to examine subgroups and interactions	Pages 5 and 6
		(c) Explain how missing data were addressed	Not applicable
		(d) If applicable, describe analytical methods taking account of sampling strategy	Not applicable
		(e) Describe any sensitivity analyses	Not applicable
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—e.g. numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	All participants eligible completed the study. Pages 6-7
		(b) Give reasons for non-participation at each stage	Not applicable
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (e.g. demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	(Not applicable)
Outcome data	15*	Report numbers of outcome events or summary measures	Page 7 and page 8
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1.
		(b) Report category boundaries when continuous variables were categorized	Not applicable
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Not applicable
Other analyses	17	Report other analyses done—e.g. analyses of subgroups and interactions, and sensitivity analyses	Figure 2. different maturity level compared.
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Page 10.
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 13.
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 13.
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 13.
<b>Other information</b>			

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Not applicable
---------	----	---	----------------

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).

# BMJ Open

## Sports Injuries Aligned to Predicted Mature Height in Highly Trained Middle-Eastern Youth Athletes: A Cohort study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-023284.R2
Article Type:	Research
Date Submitted by the Author:	11-Jan-2019
Complete List of Authors:	Rejeb, Abdallah; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie Johnson, Amanda; Aspetar Qatar Orthopaedic and Sports Medicine Hospital, Aspire Academy Sports Medicine Center Farooq, Abdulaziz; Aspetar, Orthopaedic and Sports Medicine Department, Athlete Health and Performance Research Verrelest, Ruth ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapie Pullinger, Samuel; ASPIRE Academy for Sports Excellence, Sports Sciences Vaeyens, Roel ; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Department of movement and Sports Sciences Witvrouw, Erik; Universiteit Gent Faculteit Geneeskunde en Gezondheidswetenschappen, Rehabilitation Sciences and Physiotherapy
<b>Primary Subject Heading</b>:	Sports and exercise medicine
Secondary Subject Heading:	Paediatrics, Evidence based practice
Keywords:	Youth, Biological maturation, Skeletal age, Anthropometrics, Sports injury, Mature height

SCHOLARONE™  
Manuscripts

**Sports Injuries Aligned to Predicted Mature Height in Highly Trained Middle-Eastern Youth Athletes: A Cohort study**

**Authors:**

Abdallah Rejeb<sup>1 2</sup>, Amanda Johnson<sup>1</sup>, Abdulaziz Farooq<sup>1</sup>, Ruth Verrelest<sup>2</sup>, Samuel Pullinger<sup>3</sup>, Roel Vaeyens<sup>4</sup>, Erik Witvrouw<sup>2</sup>

**Affiliations:**

<sup>1</sup> Aspetar Orthopedic Sports Medicine and Hospital, Doha, Qatar

<sup>2</sup> Ghent University, Department of Rehabilitation Sciences and Physiotherapy, Belgium

<sup>3</sup> Aspire Academy for Sports Excellence, Doha, Qatar

<sup>4</sup> Ghent University, Department of Movement and Sports Sciences, Belgium

**Corresponding author**

Abdallah Rejeb

Address: Asian Games 2006 Road, P.O. Box 22287, Aspire Academy for Sports, Doha, Qatar

Phone Number: +974 66230638

Email: [Abdallah.raheb@aspetar.com](mailto:Abdallah.raheb@aspetar.com)

Manuscript word count = 3201



**ABSTRACT** (296/300 words)

**OBJECTIVES:** To investigate the association of maturity status with injury incidence in Middle-Eastern youth athletes.

**DESIGN:** Prospective cohort study

**SETTING:** Four consecutive seasons (2010 to 2014), Aspire Academy, Qatar.

**PARICIPANTS:** Male athletes (age range: 11–18 y) representing four disciplines enrolled and grouped into two categories: individual sports and racquet sports.

**OUTCOME MEASURES:** Injury data collected over four seasons. Athletes' anthropometric characteristics assessed to calculate APHV. Predicted mature heights (PMHs) collected and categorized into four quartiles. Athletes had wrist and hand radiographs for assessment of skeletal age (SA). Early and late maturers with an SA of >1 y older or younger than their chronological age (CA).

**RESULTS:** For the sample (n = 67) across all groups, 43 (64%) athletes had one or more injuries: total of 212 injuries, 4.9 injuries per athlete across study. Survival analysis of maturity status using SA found early maturing athletes had two-fold greater injury risk compared to late maturers ([HR]; 2.04, 95% CI 1.15–3.61, P = 0.015). PMH associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006).

Athletes in 4th quartile ( $\geq 184$  cm) had up to 2-fold injury risk (HR 2.41, 95% CI 1.42–4.08, P = 0.001). Racquet and individual sports involved similar injury risk (HR 1.14, 95% CI 0.86–1.52, P = 0.37).

**CONCLUSION:** SA early maturity and PMH gradient were significant predictors of injury in youths.

**STRENGTHS AND LIMITATIONS OF THE STUDY**

- First longitudinal study to assess anthropometric characteristics and biological maturity status as injury risk factors in Middle-Eastern athletes.
- Participants were highly trained adolescent athletes.
- Measurement of maturity and growth were at moderate-to-high risk of bias.
- SA has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment and interpretation of results.

**INTRODUCTION**

The range of somatic and biological maturity in individuals of the same chronological age (CA) is large [1]. Such observations are derived from correlational and multivariate studies that compare young individuals of the same age who are at both extremes of the maturity range [2]. Therefore, the assessment of maturity is an important consideration when dealing with adolescent athletes on a longitudinal basis. Further, understanding the cause of disease and injury is vital in predicting and preventing injury [3].

In young athletes, the demands of their chosen sport are superimposed on normal growth and maturation. A literature review revealed that there is a greater susceptibility to injury during certain periods of growth [4–6]. Indeed, the association between an increased prevalence of injuries and the adolescent growth spurt has long been recognized [7–9]. A recent study analysis [10] on adolescent soccer players revealed greater risk of injury with players within age at peak height velocity (APHV) in comparison with the players before and after APHV. Mismatched rapid growth in the long bones relative to muscular lengthening may disrupt structure, neuromuscular function, and physical performance [11].

Deehan et al. [12], state that an increased participation in sports predisposes the immature skeleton to injury. Furthermore, participation in high intensity sport entails an inherent risk of

sports-related injuries, and this is heightened at various stages of growth and maturation [13]. Maturation induces profound changes in the skeletal, neuromuscular, and tendinous systems of young athletes [14] and mismatches in biological maturity may create competitive inequality and increase the risk of injury [15]. Le Gall et al. [16], further point out that injury rates generally increase with increasing CA. However, CA is a poor indicator of biological maturity [17]; moreover, Ardern et al. [18], report that chronological age alone is an unreliable indicator of skeletal maturity. Skeletal age (SA) is generally accepted as the most accurate method of assessing biological maturity [6,19], by identifying critical periods of development; it also offers a rational method for monitored age-specific training. Before initiating any program for mitigating sports injuries, the magnitude of the problem must be identified and the extent of the injury defined in terms of incidence and severity [20].

A number of studies have been conducted involving injuries in adolescent footballers; conversely, few studies have focused on injuries in non-footballer adolescent athletes in high performance sporting environments [21]. Studies of anthropometric characteristics and biological maturity status as injury risk factors in Middle-Eastern youths are also limited, highlighting the need for more research in this area. Therefore, the purpose of the present study was to investigate injury incidence according to biological maturity using two outcome measures (SA and PHV) in highly trained youth athletes based at a Middle Eastern Sports Academy.

## METHODS

Sixty-seven highly trained adolescent athletes (age range 11–18 y) representing athletics and racquet sports (table tennis and squash) from a Middle Eastern sports school were included in this four-year study. A prospective, longitudinal cohort design was used and included separate observation periods over four consecutive seasons (2010–2011, 2011–2012, 2012–2013, and 2013–2014), i.e., school years, which lasted from the beginning of September until the end of June (~40 weeks). Participant maturity assessments included both anthropometric

measurements, collected three times a season, and SA assessments using Fels method completed once, at the start of every season. Medical screening was performed at the beginning of each season to determine health and injury status. All selected athletes had clearance from a physician to participate in their respective sport. Written informed consent was sought and obtained from parents and assent from all participants. The study was part of a general sports science provision to the sports academy, and all procedures were reviewed and granted by the Institutional Review Board (IRB) for Human Subjects and conformed to the recommendations of the Declaration of Helsinki.

**Participants**

Figure 1 shows the flow of participants in the study over consecutive seasons. A total of four sporting disciplines were analysed, grouped into two categories: athletics and fencing and racquet sports (squash and table tennis). This classification was based on specific sport characteristics and injury risk [22,23]. Inclusion criteria were as follows: (1) the athlete had to be enrolled in the sports school during at least one full school year; (2) athletes with injuries in previous seasons were not excluded from this study, but injuries present at the beginning of the observation period were not included in statistical analyses; and (3) injuries that were not sustained in the context of the sports program (e.g. recreational activities) or data related to sickness or other general medical conditions were not used for further analysis.

<<<Insert Figure 1 here>>>

**Injury definition and data collection**

An Injury was defined as any physical complaint, which occurred during sports training, strength and conditioning training or during competition. Injuries were divided into time-loss (TL) injuries and no time-loss (NTL) injuries. A clinical examination and/or treatment of an athlete which did

not result in a full training session or competition being missed was described as a complaint with NTL injury. A clinical examination and/or treatment of an athlete resulting in a training session or competition being missed the following day(s) was labelled as a TL injury [23]. A traumatic injury was defined as any injury resulting from a specific and identifiable mechanism, including contact and non-contact circumstances with acute onset. Overuse injuries were defined as injuries resulting from insidious onset without a recognizable mechanism. Injury severity was defined, based on days of absence from usual sport participation, as slight (1 d or less), minimal (2–3 d), mild (4–7 d), moderately serious (8–28 d), serious (>28 d up to 6 months) or long-term (>6 months) in accordance with [24].

All injuries were collected by a physical therapist (AR) with experience of working within youth sport. Data from medical records were used to document all sports related injuries during the study. Each sporting discipline had a dedicated full-time physiotherapist and a full-time employed medical doctor at the sports academy. The medical record used an injury reporting system based upon the football injury reporting system [25] and the Sport Medicine Diagnostic Coding System [26]. Information was gathered concerning all injuries related to sports activity, including several related variables (e.g. type, location, affected structure, mechanism [acute vs. overuse], time loss, severity, and date of injury).

### **Somatic maturation and anthropometric measurements**

Anthropometric measurements were initially carried out on all participants on a three monthly basis along with an estimation of the age at peak height velocity (APHV) as a relative indicator of somatic maturity and representing the time of maximum growth in stature during adolescence using Mirwald method [27] for the prediction of growth [1]. APHV was calculated from the first measurement recorded. To ensure that the outcome measures remained consistent and reliable, every effort was made to ensure that measurements were taken at approximately the same time of the season. Measurements were collected by qualified practitioners from the International

Society for the Advancement of Kinanthropometry (ISAK) and included stretch stature ( $\pm 0.1$  cm Holtain Limited, Crosswell, UK).

The predicted mature height (PMH) of all participants were collected and categorized into four PMH quartiles (Q1–Q4: Q1 < 176 cm; 176 cm  $\leq$  Q2 < 180 cm; 180 cm  $\leq$  Q3 < 184 cm; Q4  $\geq$  184 cm). The athletes were then divided into three maturity groups (late, normal, or early maturing) based on the mean  $\pm 1.0$  year of the APHV of the total sample (late, APHV > mean + 1.0 y; normal, APHV within mean  $\pm 1.0$  y; early, APHV < mean – 1.0 y). Years from peak height velocity (maturity offset value: CA – maturity offset) was calculated by subtracting the CA at the date of injury from the age at estimated peak height velocity.

**Skeletal maturation assessment**

Each year athletes were required to have a radiograph of the left wrist and hand, a convenient area to examine, and a more accurate method for the assessment of SA [11], using the Fels method [6,28] which has an advantage over other methods [29]. Maturity status, defined by the difference between CA and SA was calculated and classified into four categories: late, normal, early, and mature athletes. Late referred to an SA that was younger than CA by more than 1.0 y, athletes with a normal pattern of maturity had an SA that was within 1.0 y of CA, early referred to an SA that was older than CA by more than 1.0 y, and the closure of growth plate determine skeletally mature athletes.

**Statistical Analysis**

Data were analysed using statistical software (Stata Corporation, College Station, Texas). Descriptive statistics were presented as frequencies and proportions (%), and incidence rates were expressed as the number of injuries/number of registered athletes. To examine the role of growth status and maturity with the onset of injuries, a univariate Cox regression survival analysis was performed after accounting for repeated visits of athletes over the four seasons. Hazard ratios

(HR) with 95% confidence intervals (CIs) were reported for each factor. Kaplan-Meier curves were plotted for SA groups and time to injury over a season. Where appropriate, 95% CIs are presented. The alpha level of significance was set at 5%.

### Patient and public involvement statement

Patients and public were not involved in the analysis of this study.

## RESULTS

Throughout the four-year seasons study period, 67 athletes were enrolled representing 151 athletic seasons. Table 1 presents the anthropometric characteristics of participants and their maturity status. From these participants, 43 (64%) reported one or more injuries adding up to 212 injuries in total. The injury rate observed per registered athlete amounted to 4.9 injuries over the course of four seasons.

**Table 1.** Anthropometric characteristics (mean  $\pm$  SD) of participants according to maturity status

	Late	Normal	Early
	(n = 4, 6.0%)	(n = 59, 88.1%)	(n = 4, 6.0%)
CA (years)	13.3 $\pm$ 1.1	12.3 $\pm$ 1.0	12.1 $\pm$ 0.5
Years from PHV	-2.4 $\pm$ 1.2	-1.6 $\pm$ 1.1	-0.1 $\pm$ 0.9
APHV (years)	15.8 $\pm$ 1.5	13.9 $\pm$ 0.5	12.2 $\pm$ 0.9
PMH (cm)	181.6 $\pm$ 17.1	179.4 $\pm$ 4.9	188.4 $\pm$ 3.5
% PMH (%)	85.0 $\pm$ 3.0	85.0 $\pm$ 4.0	90.0 $\pm$ 4.0
SA (years)	11.8 $\pm$ 0.5	12.8 $\pm$ 1.5	12.7 $\pm$ 1.8

CA: chronological age; APHV: age at peak height velocity; PMH: predicted mature height; SA: skeletal age; SD: standard deviation

### Skeletal age: maturity status distribution and injury risk

Among all participants (n = 67), 4% were classified as late maturers, 33% as normal, 41% as early and 22% as skeletally mature. The overall injury free survival analysis of maturity status using SA assessment indicated that early maturing athletes had a two-fold higher risk of injury over a season compared to late maturing athletes (HR 2.04, 95% CI 1.15–3.61, P = 0.015),

(Figure 2). There was a trend that early maturing athletes had a greater risk of injury over a season compared to normal athletes (HR 1.62, 95% CI 0.99–2.65,  $P = 0.053$ ), but this was only marginally significant. However, injury risk among late and fully mature athletes did not differ from normal maturers.

<<<<Insert Figure 2 Here >>>>

**Somatic maturation and anthropometric measurements: distribution and injury risk**

Using anthropometric measurements, among all participants ( $n = 67$ ), 6.0% were classified as late maturing, 85.8% as normal, and 6.0% as early. Classification of participant maturity status (late, normal, and early) according to age at PHV (APHV) was not significantly associated with overall injury incidence in this cohort of highly trained Middle-Eastern youth athletes. Older PHVs were marginally associated with higher injury risk, but this was not statistically significant (HR 1.11, 95% CI 0.99–1.23,  $P = 0.067$ ).

Both PMH (cm), and %PMH were found to be associated with injury risk (HR 1.05, 95% CI 1.01–1.08,  $P = 0.006$ , and HR 1.03, 95% CI 1.00–1.06,  $P = 0.026$ ), respectively. When compared to participants in the 1st quartile for PMH ( $<176$ ), athletes in the 4th quartile ( $\geq 184$  cm) had a two and half times greater risk of injury (HR 2.41, 95% CI 1.42–4.08,  $P = 0.001$ ) over a season.

No significant differences were observed in the injury risk between racquet sports ( $n = 30$ ) and individual sports athletes ( $n = 37$ ; HR 1.14, 95% CI, 0.86–1.52,  $P = 0.37$ ).

**DISCUSSION**

The present investigation was carried out to examine injury incidence according to maturity status. Biological maturity status and height gradient play a significant role in injury risk profiles of highly trained youth athletes. The results of the current study show that athletes maturing at a younger



age are at significantly greater risk of injury, more than two-fold, compared to their later maturing counterparts. Taller athletes were also found to be significantly more at risk of injury.

There is limited and contrasting evidence on the relationship between maturity and injury in youth sports [10,30,31]. In this study, SA maturity (Fels method) showed that early maturing athletes had twice the risk of injury over a season compared to late maturing athletes. This finding is consistent with previous study [6], that described that early maturing athletes are significantly more at risk of injury than late or normally maturing athletes. A possible explanation could be that youth players with higher engagement and performance advantages are often associated with early maturation, usually transient during adolescence, and maybe reversed in early adulthood [16]

However, our study results were inconsistent with other study [30] on youth athletes, in which late maturing athletes have a higher injury rate compared to their earlier maturing counterparts. A plausible explanation could be that Fourchet et al. [30], examined anthropometric data collected from a track and field cohort for their findings, while our study resulted from maturity status derived from bone age but with no substantial association from APHV.

In the present study, no significant association was observed between APHV and injury risk (HR 0.90, 95% CI 0.74–1.11,  $P = 0.329$ ), which is inconsistent with recent data on youth alpine ski racing [32] and other studies on talented Dutch and English youth soccer players [6,33] which show a heightened period of risk around the time of peak height velocity. An explanation of these discrepancies could be that our study cohort was not large enough, as the APHV method appears to be useful in youth talent selection and injury prevention programs because it can be easily applied in a large cohort of young athletes [34].

PMH and %PMH at a given age are minimally invasive, feasibly practical indicators of somatic maturation [17,35], especially if mature height can be assessed without an estimate of SA [28]. In

this study, the PMH and %PMH revealed that both indicators were associated with injury risk (HR 1.05, 95% CI 1.01–1.08, P = 0.006), and HR 1.03, 95% CI 1.00–1.06, P = 0.026), respectively. When compared to participants in the 1st quartile for PMH (<176), athletes in the 4th quartile ( $\geq 184$  cm) had two and a half times greater risk of injury (HR 2.41, 95% CI 1.42–4.08, P = 0.001). The present results are partly in line with previous studies on other sports. Johnson et al. [6], showed that gains in height in youth footballers over a season were associated with an increased number of injuries. The study of Kemper et al. [36], on elite youth soccer players with growth rates of at least 0.6 cm/month showed a higher risk for injury. In a different study on soccer athletes, it was found that the tallest boys had the highest incidence of injury [37]. However, these findings and those of the present study are not in line with a study on youth football players [38], in which injured and non-injured players did not differ in percentage of mature height. An explanation could be that the definition of reportable injury in the methods of the study, which considered only time loss injuries, did not capture the full spectrum of injuries and therefore overlooked other injuries with insidious onset e.g. growth conditions.

The results of this study have some important practical implications. Malina et al. [2], advocate the documentation of anthropometric characteristics, biological maturity, and physical fitness parameters as crucial aids in the prevention of injury. Noninvasive methods for estimating maturity status may allow youth programs to match players using maturity status rather than CA, and thus equalize competition to some extent. An unequal competition is regarded as an impediment to personal development [39]. Furthermore, it has been suggested that there is an overwhelming bias in sport favoring taller athletes [40], and data on Olympic medal winners show that many running and jumping events are seriously biased in favor of the very tall [41].

When examining the classification resulting from SA of late (4%), normal (33%), early (41%), and skeletally mature athletes (22%), the under-representation of late and preponderance of early maturing athletes in this cohort is consistent with observations for male youth athletes in several

sports including soccer and alpine ski racing [10,19,32]. However, these results and those of the present study are not in line with the study of Johnson et al. [6], on schoolboy footballers, in which two thirds of their players fall within the normal maturity category. Moreover, Le Gall et al. [16], classify only 12.0% as late maturers, 63.5% as normal maturers, and 24.5% as early maturers. These discrepancies are believed to be due to differences in selection policies and talent identification policies (physical, technical, and tactical skills) of varying elite development centers. Several studies point out that athletes who are more advanced in their biological maturity perform better than their later maturing peers and have a better chance of being selected [42–44]. Youth sport is highly selective, with a maturity-associated selection/exclusion process [35].

### **Implications and concepts for prevention**

The findings in this study have several implications for youth athletes. First, our data suggests that adolescent athletes might be identified and selected with a preference for youths with advanced maturity. Such selection strategies which favor early maturers entail significant risks of injury. Accordingly, those involved in the selection and development of young athletes should be cognizant of temporary changes in motor control that may occur during these periods [45], consider maturity status, develop appropriate training programs to optimize training adaptation, design injury prevention plans to minimize activity related injury risk, and mitigate long term youth injury consequences.

Limitations of the current study should be noted. First, biological maturation methods have inherent limitations when applied to youth athletes and need to be applied with caution. Although SA is a gold standard indicator of maturation, it has major limitations in expense and minimal radiation and lack of knowledgeable staff for assessment protocols and the interpretation of results [46]. Although our sample size is small, we have a follow-up over four seasons. Another limitation, we had no data on training or competition exposure, which reduces the comparability with other studies reporting injury incidence.

It must also be remembered that, except for accidents, a sports injury can rarely be ascribed to a single factor, but rather to an association of causes or circumstances and the interaction among a web of determinants [47,48].

CONCLUSIONS

The findings of the present study showed that maturity status plus PMH and %PMH are associated with injury in individual and racquet sports but no association has been established between APHV and injury. As SA varies individually in rate and timing, and mismatches in maturity may create competitive inequality and increase injury incidence, it is suggested that biological maturity should be considered during training to help prevent injury. Given the peculiarity of youth athletes it is important to optimize the planning of training activities to further improve the understanding of the link between training, growth, and injury.

Figure legends

<<<Figure 1. Flowchart describing the inclusion and flow of participants throughout the study >>>  
<<<Figure 2. Kaplan-Meier survival analysis of injuries in relation to different skeletal age (SA) maturity status >>>

REFERENCES

1 Iuliano-Burns S, Mirwald RL, Bailey DA. Timing and magnitude of peak height velocity and peak tissue velocities for early, average, and late maturing boys and girls. *Am J Hum Biol* 2001;**13**:1–8. doi:10.1002/1520-6300(200101/02)13:1<1::AID-AJHB1000>3.0.CO;2-S

2 Malina RM, Cumming SP, Morano PJ, et al. Maturity status of youth football players: A noninvasive estimate. *Med Sci Sports Exerc* 2005;**37**:1044–52. doi:10.1249/01.mss.0000171622.45134.cb

- 3 Meeuwisse W. Assessing Causation in Sport Injury: A Multifactorial Model. *Clin J Sport Med* 1994;**4**:166–70. doi:10.1097/00042752-199407000-00004
- 4 Krivickas LS, Feinberg JH. Lower extremity injuries in college athletes: Relation between ligamentous laxity and lower extremity muscle tightness. *Arch Phys Med Rehabil* 1996;**77**:1139–43. doi:10.1016/S0003-9993(96)90137-9
- 5 Aicardi G, Vignolo M, Milani S, *et al.* Assessment of skeletal maturity of the hand-wrist and knee: A comparison among methods. *Am J Hum Biol* 2000;**12**:610–5. doi:10.1002/1520-6300(200009/10)12:5<610::AID-AJHB5>3.0.CO;2-D
- 6 Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. *Bmj* 2009;**338**:b490–b490. doi:10.1136/bmj.b490
- 7 Peterson HA. *Epiphyseal growth plate fractures*. 2007. doi:10.1007/978-3-540-33802-4
- 8 Bailey DA, Wedge JH, McCulloch RG, *et al.* Epidemiology of fractures of the distal end of the radius in children as associated with growth. *J Bone Jt SurgAm* 1989;**71**:1225–31. <http://www.ncbi.nlm.nih.gov/pubmed/2777851>
- 9 Leppänen M, Pasanen K, Clarsen B, *et al.* Overuse injuries are prevalent in children's competitive football: a prospective study using the OSTRC Overuse Injury Questionnaire. *Br J Sports Med* 2018;:bjsports-2018-099218. doi:10.1136/bjsports-2018-099218
- 10 Materne O, Farooq A, Johnson A, *et al.* Relationship between injuries and somatic maturation in highly trained youth soccer players. *Int Res Sci Soccer II* 2016;:182–92. <https://books.google.com/books?hl=en&lr=&id=w8DMCgAAQBAJ&oi=fnd&pg=PA182&dq=Relationship+between+injuries+and+somatic+maturation+in+highly+trained+youth+soccer+players&ots=6D7Xy7rLzj&sig=ywUDNrV1u4pMrJIBN3wERj1u93M%5Cnhttp://s3>

amazonaws.com/academia.edu.

11 Lloyd RS, Oliver JL, Faigenbaum AD, *et al.* Chronological Age vs. Biological Maturation. *J Strength Cond Res* 2014;**28**:1454–64. doi:10.1519/JSC.0000000000000391

12 Deehan DJ, Bell K, McCaskie AW. Adolescent musculoskeletal injuries in a football academy. *J Bone Jt Surg - Br Vol* 2007;**89-B**:5–8. doi:10.1302/0301-620X.89B1.18427

13 Read P, Oliver JL, De Ste Croix MBA, *et al.* Injury Risk Factors in Male Youth Soccer Players. *Strength Cond J* 2015;**37**:1–7. doi:10.1519/SSC.0000000000000171

14 Huey K, Robinson MM, Wiesinger H-P, *et al.* Imbalances in the Development of Muscle and Tendon as Risk Factor for Tendinopathies in Youth Athletes: A Review of Current Evidence and Concepts of Prevention. 2017;**8**:1–18. doi:10.3389/fphys.2017.00987

15 Cumming SP, Brown DJ, Mitchell S, *et al.* Premier League academy soccer players' experiences of competing in a tournament bio-banded for biological maturation. *J. Sports Sci.* 2017;;1–9. doi:10.1080/02640414.2017.1340656

16 Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. *Scand J Med Sci Sport* 2007;**17**:564–72. doi:10.1111/j.1600-0838.2006.00594.x

17 Beunen GP, Rogol AD, Malina RM. Indicators of biological maturation and secular changes in biological maturation. *Food Nutr. Bull.* 2006;**27**. doi:10.1177/156482650903000307

18 Ardern CL, Ekås G, Grindem H, *et al.* 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Knee Surgery, Sport Traumatol Arthrosc* 2018;**0**:1–17. doi:10.1007/s00167-018-4865-y

19 Malina RM. Early sport specialization: Roots, effectiveness, risks. *Curr Sports Med Rep*

- 2010;**9**:364–71. doi:10.1249/JSR.0b013e3181fe3166
- 20 Leppänen M, Lapinleimu H, Lehtonen L, *et al*. Growth of extremely preterm infants born in 2001-2010. *Acta Paediatr Int J Paediatr* 2013;**102**:206–8. doi:10.1111/apa.12061
- 21 Steffen K. More data needed on injury risk among young elite athletes. *Br J Sports Med* 2010;**44**:485–
- 9.http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=51172712&site=ehost-live
- 22 Rejeb A, Johnson A, Vaeyens R, *et al*. Compelling overuse injury incidence in youth multisport athletes. *Eur J Sport Sci* 2017;**17**:495–502.
- doi:10.1080/17461391.2016.1275820
- 23 Malisoux L, Frisch A, Urhausen A, *et al*. Injury incidence in a sports school during a 3-year follow-up. *Knee Surgery, Sport Traumatol Arthrosc* 2013;**21**:2895–900.
- doi:10.1007/s00167-012-2185-1
- 24 Timpka T, Alonso J-M, Jacobsson J, *et al*. Injury and illness definitions and data collection procedures for use in epidemiological studies in Athletics (track and field): Consensus statement. *Br J Sports Med* 2014;**48**:483–90. doi:10.1136/bjsports-2013-093241
- 25 Fuller CW, Ekstrand J, Junge A, *et al*. Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *Scand. J. Med. Sci. Sport*. 2006;**16**:83–92. doi:10.1111/j.1600-0838.2006.00528.x
- 26 Meeuwisse W, H MD P, Tyreman H, *et al*. A Dynamic Model of Etiology in Sport Injury: The Recursive Nature of Risk and Causation. *Clin J Sport Med* 2007;**17**:215–9.
- doi:10.1097/JSM.0b013e3180592a48

- 1  
2  
3 27 MIRWALD RL, G. BAXTER-JONES AD, BAILEY DA, *et al.* An assessment of maturity  
4 from anthropometric measurements. *Med Sci Sport Exerc* 2002;**34**:689–94.  
5  
6 doi:10.1097/00005768-200204000-00020  
7  
8  
9  
10 28 Khamis HJ, Roche AF. Predicting adult stature without using skeletal age: the Khamis-  
11 Roche method. *Pediatrics*. 1994;**94**:504–7. doi:10.1016/S0740-624X(98)90020-X  
12  
13  
14  
15 29 Silventoinen K, Pke M, Tynelius P. Muscle Strength and Body Size and Later  
16 Cerebrovascular and Coronary Heart Disease Maturity Status and Injury Risk in Youth  
17 Soccer Players. *Sport Med* 2010;**20**:131–5. doi:10.1136/bmj.b490.Objective  
18  
19  
20  
21  
22 30 Fourchet F, Horobeanu C, Loepelt H, *et al.* Foot, Ankle, and Lower Leg Injuries in Young  
23 Male Track and Field Athletes. *Int J Athl Ther Train* 2011;**16**:19–23 5p.  
24  
25 doi:10.1123/ijatt.16.3.19  
26  
27  
28  
29 31 Swain M, Kamper SJ, Maher CG, *et al.* Relationship between growth, maturation and  
30 musculoskeletal conditions in adolescents: a systematic review. *Br J Sports Med*  
31 2018;**bjsports-2017-098418**. doi:10.1136/bjsports-2017-098418  
32  
33  
34  
35  
36 32 Müller L, Hildebrandt C, Müller E, *et al.* Long-Term athletic development in youth alpine  
37 ski racing: The effect of physical fitness, ski racing technique, anthropometrics and  
38 biological maturity status on injuries. *Front Physiol* 2017;**8**. doi:10.3389/fphys.2017.00656  
39  
40  
41  
42  
43 33 Vanderlei FM, Vanderlei LCM, Bastos FN, *et al.* Characteristics and associated factors  
44 with sports injuries among children and adolescents. *Brazilian J Phys Ther* 2014;**18**:530–  
45 7. doi:10.1590/bjpt-rbf.2014.0059  
46  
47  
48  
49  
50 34 Müller L. Müller EHCKK& RC. Die Erhebung des biologischen Entwicklungsstandes für  
51 die Talentselektion - welche Methode eignet sich? [The assessment of biological  
52 maturation for talent selection - which method can be used?]. *Sport Sport* 2015;**29**:56–63.  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 35 Bergeron MF, Mountjoy M, Armstrong N, *et al.* International Olympic Committee  
4 consensus statement on youth athletic development. *Br J Sports Med* 2015;**49**:843–51.  
5 doi:10.1136/bjsports-2015-094962  
6  
7  
8  
9  
10 36 Kemper GLJ, Van Der Sluis A, Brink MS, *et al.* Anthropometric Injury Risk Factors in  
11 Elite-standard Youth Soccer. *Int J Sports Med* 2015;**36**:1112–7. doi:10.1055/s-0035-  
12 1555778  
13  
14  
15  
16  
17 37 Backous DD, Friedl KE, Smith NJ, *et al.* Soccer Injuries and Their Relation to Physical  
18 Maturity. *Am J Dis Child* 1988;**142**:839–42. doi:10.1001/archpedi.1988.02150080045019  
19  
20  
21  
22 38 Malina RM, Morano PJ, Barron M, *et al.* Incidence and player risk factors for injury in  
23 youth football. *Clin J Sport Med* 2006;**16**:214–22. doi:10.1097/00042752-200605000-  
24 00005  
25  
26  
27  
28  
29 39 Musch J, Grondin S. Unequal Competition as an Impediment to Personal Development: A  
30 Review of the Relative Age Effect in Sport. *Dev Rev* 2001;**21**:147–67.  
31 doi:10.1006/drev.2000.0516  
32  
33  
34  
35  
36 40 Baxter-Jones ADG. Growth and Development of Young Athletes: Should Competition  
37 Levels be Age Related? *Sport Med* 1995;**20**:59–64. doi:10.2165/00007256-199520020-  
38 00001  
39  
40  
41  
42  
43 41 Khosla T. Unfairness of certain events in the Olympic games. *Br Med J* 1968;**4**:111–3.  
44 doi:10.1136/bmj.4.5623.111  
45  
46  
47  
48 42 Beunen GP, Malina RM, Van't Hof MA, *et al.* *Adolescent Growth and Motor Performance.*  
49 1988.  
50  
51  
52  
53 43 Philippaerts RM, Vaeyens R, Janssens M, *et al.* The relationship between peak height  
54 velocity and physical performance in youth soccer players. *J Sports Sci* 2006;**24**:221–30.  
55  
56  
57  
58  
59  
60

doi:10.1080/02640410500189371

44 Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity. *Growth, Matur Phys Perform* 2004;:1–17.

45 Mueller L, Hildebrandt C, Mueller E, *et al.* Injuries and illnesses in a cohort of elite youth alpine ski racers and the influence of biological maturity and relative age: a two-season prospective study. *Open Access J Sport Med* 2017;**Volume 8**:113–22.  
doi:10.2147/OAJSM.S133811

46 Malina RM, Rogol AD, Cumming SP, *et al.* Biological maturation of youth athletes: assessment and implications. *Br J Sports Med* 2015;**49**:852–9. doi:10.1136/bjsports-2015-094623

47 Theisen D, Malisoux L, Seil R, *et al.* : Injuries in Youth Sports: Epidemiology, Risk Factors and Prevention. *Dtsch Z Sportmed* 2014;**65**:S248-252.<http://www.zeitschrift-sportmedizin.de/artikel-online/archiv-2014/heft-9/injuries-in-youth-sports-epidemiology-risk-factors-and-prevention/>

48 Bittencourt NFN, Meeuwisse WH, Mendonça LD, *et al.* Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition—narrative review and new concept. *Br J Sports Med* 2016;**50**:1309–14.  
doi:10.1136/bjsports-2015-095850

**CONTRIBUTORSHIP STATEMENT:** AR designed and developed the research question and wrote the original version of the manuscript as part of his doctoral thesis. EW (doctoral supervisor) reviewed, designed and provided expertise to the study. AJ (doctoral committee member) was involved in study design. AF supervised and provided expertise with respect to the data analyses. RVe, SP, and RVa (doctoral committee member) reviewed and provided expertise to the study. All authors have contributed to and edited the manuscript and have approved the final manuscript.

**FUNDING:** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

**COMPETING INTERESTS:** None declared.

**ETHICS APPROVAL:** The study received ethical approval by the Anti- Doping Lab Qatar

**DATA SHARING STATEMENT:** No additional data are available

**ACKNOWLEDGEMENTS:** The publication of this article was funded by the Qatar National Library.




Figure 1. Flowchart describing the inclusion and flow of participants throughout the study  
135x89mm (300 x 300 DPI)




Figure 2. Kaplan-Meier survival analysis of injuries in relation to different skeletal age (SA) maturity status

96x67mm (300 x 300 DPI)

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cross-sectional studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	Page 1 and Page 2.
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Pages 2 and 3.
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Pages 4.
Objectives	3	State specific objectives, including any prespecified hypotheses	Pages 4 and 5.
Methods			
Study design	4	Present key elements of study design early in the paper	Page 5, line of methods 86
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Page 5.
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Pages 5 and 6. Figure 1.
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	On pages 6, 7 and 8. Definition and data collection of outcome variables were given.
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	On Pages 7 and 8.
Bias	9	Describe any efforts to address potential sources of bias	On Page 8.
Study size	10	Explain how the study size was arrived at	Every available athlete was included in our study.
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Page 5.

Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	Pages 8.
		(b) Describe any methods used to examine subgroups and interactions	Pages 5 and 6
		(c) Explain how missing data were addressed	Not applicable
		(d) If applicable, describe analytical methods taking account of sampling strategy	Not applicable
		(e) Describe any sensitivity analyses	Not applicable
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—e.g. numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	All participants eligible completed the study. Pages 6-7
		(b) Give reasons for non-participation at each stage	Not applicable
		(c) Consider use of a flow diagram	Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (e.g. demographic, clinical, social) and information on exposures and potential confounders	Table 1
		(b) Indicate number of participants with missing data for each variable of interest	(Not applicable)
Outcome data	15*	Report numbers of outcome events or summary measures	Page 7 and page 8
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (e.g., 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Table 1.
		(b) Report category boundaries when continuous variables were categorized	Not applicable
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Not applicable
Other analyses	17	Report other analyses done—e.g. analyses of subgroups and interactions, and sensitivity analyses	Figure 2. different maturity level compared.
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	Pages 8-10.
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Page 13.
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Page 13.
Generalisability	21	Discuss the generalisability (external validity) of the study results	Page 13.
<b>Other information</b>			

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Not applicable
---------	----	---	----------------

\*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at [www.strobe-statement.org](http://www.strobe-statement.org).